arXiv:hep-th/0311186v4 16 Mar 2004

hep-th/0311186
HU-EP 03/181

Type IIA on a Compact Calabi-Yau and $D = 11$ Supergravity Uplift of its Orientifold

Aalok Misra[1]
Institut für Physik, Humboldt Universität, Berlin,
Germany

Abstract

Using the prescription of [1] for defining period integrals in the Landau-Ginsburg theory for compact Calabi-Yau's, we obtain the Picard-Fuchs equation and the Meijer basis of solutions for the compact Calabi-Yau $CY_3(3, 243)$ expressed as a degree-24 Fermat hypersurface *after* resolution of the orbifold singularities. This is similar in spirit to the method of obtaining Meijer basis of solutions in [2] for the case wherein one is away from the orbifold singularities, and one is considering the large-base limit of the Calabi-Yau. The importance of the method lies in the ease with which one can consider the large *and* small complex structure limits, as well as the ability to get the "ln"-terms in the periods without having to parametrically differentiate infinite series. We consider in detail the evaluation of the monodromy matrix in the large and small complex structure limits. We also consider the action of the freely acting antiholomorphic involution of [3, 2] on $D = 11$ supergravity compactified on $CY_3(3, 243) \times S^1$ [4] and obtain the Kähler potential for the same in the limit of large volume of the Calabi-Yau. As a by-product, we also give a conjecture for the action of the orientation-reversing antiholomorphic involution on the periods, given its action on the cohomology, using a canonical (co)homology basis. Finally, we also consider showing a null superpotential on the orientifold of type IIA on $CY_3(3, 243)$, having taken care of the orbifold singularities, thereby completing the argument initiated in [2].

---

[1] Address from Jan 3, 2004: Indian Institute of Technology, Roorkee, India

1 Introduction

The periods are the building blocks, e.g., for getting the prepotential (and hence the Kähler potential and the Yukawa coupling) in $\mathcal{N} = 2$ type II theories compactified on a Calabi-Yau, and the superpotential for $\mathcal{N} = 1$ type II compactifications in the rpesence of (RR) fluxes. It is in this regard that the Picard-Fuchs equation satisfied by the periods, become quite important. Also, traversing non-trivial loops in the complex structure moduli space of type IIB on a Calabi-Yau mirror to the one on the type IIA side, corresponds to shifting of the Kähler moduli in the Kähler moduli space on the type IIA side. This results in mixing of flux numbers corresponding to RR fluxes on the type IIA side, implying thereby that dimensions of cycles on the type IIA side, loose their meaning. The mixing matrix for the flux numbers is given by the monodromy matrix. The mixing matrix for the flux numbers if given by the monodromy matrix. It hence becomes important to evaluate the same. Non-compact Calabi-Yau's have typically been the ones that have been extensively studied in this context as well as topological strings with the same as target spaces. Compact Calabi-Yau's, in addition to the quintic, thus become quite important to be studied. Additionally, mirror symmetry becomes quite indispensible when working with compact Calabi-Yau's with large value of $h^{2,1}$ and small $h^{1,1}$. One such compact Calabi-Yau that we will study is $CY_3(3, 243)$, i.e., one for which $h^{1,1} = 3$, and $h^{2,1} = 243$. This compact Calabi-Yau plays a central role in $\mathcal{N} = 2, 1$ type IIA/Heterotic dual paris in four dimensions in [5, 6].

In [2], we addressed the issue of deriving the Picard-Fuchs equation on the mirror Landau-Ginsburg side corresponding to the gauged linear sigma model for a compact Calabi-Yau $CY_3(3, 243)$, expressed as a degree-24 Fermat hypersurface in a suitableweighted complex projective space, but staying away from the orbifold singularities by taking the large-base limit of the compact Calabi-Yau. Even though, one ended up with more than the required number of solutions, but the essential idea that was highlighted was the ease with which, both the large and small complex structure limits could be addressed, and the fact that the nonanalytic ln-terms in the periods, could be easily obtained without having to resort to parametric differentiations of infinite series. In this paper, we address the problem of getting the right number of the right kind of solutions on the mirror Landau-Ginsburg side, but this time after having resolved the orbifold singularities. We also address the problems of showing that unoriented instantons do not generate a superpotential on the type IIA side in the $\mathcal{N} = 1$ Heterotic/type IIA dual pair of [5], whose M and F theory uplifts were discussed in [3]. It was shown in [2], using mirror symmetry, that as expected from the Heterotic and F theory duals, there is indeed no superpotential generated from $\mathbf{RP}^2$-instantons in the type IIA side in the large-base limit of $CY_3(3,243)$, away from the aforementioned orbifold singularities of the relevant Fermat hypersurface. In this paper, we show that the same remains true even after the resolution of the orbifold singularities. Further, we discuss the supergravity uplift of the type IIA orientifold that figures in the abovementioned $\mathcal{N}=1$ Heterotic/type IIA dual pair, to $D=11$ supergravity. We evaluate the Kähler potential in the large volume limit of $CY_3(3,243)$. As an interesting aside, we give a conjecture about the action of the antiholomorphic involution that figures in the definition of the type IIA orientifold, on the periods, given its action on the cohomology of $CY_3(3,243)$, using a canonical (co)homology basis to expand the holomorphic 3-form. We verify the conjecture for T^6 and (partly) for the mirror to the quintic.

The plan of the paper is the following. In section **2**, after a brief discussion of the orbifold singularities' resolution corresponding to the Fermat hypersurface, we define the period integral as per the prescription of Hori and Vafa, for a compact Calabi-Yau. We then obtain the degree-8 Picard-Fuchs satisfied by the period integral, and then discuss the guiding principle in obtaining the right number of the right kind of solutions, and obtain a set of solutions to the equation, and evaluate the relevant contour integrals. We also discuss the evaluation of the monodromy matrix in the large and small complex structure limits. In section **3**, we then discuss showing that there is no superpotential generated (by the only possible source - unoriented instantons) for type IIA on an orientifold of $CY_3(3,243)$, *after* resolution of the aforementioned orbifold singularities, thereby strengthening the conjectured $\mathcal{N}=1$ triality in [3]. In section **4**, we discuss the $D=11$ supergravity uplift of the aforementioned type IIA orientifold on a 'barely' G_2 manifold [3], and also evaluate the Kähler potential in the limit of the large volume of the Calabi-Yau. Finally, in section **5**, we discuss a conjecture related to the action of antiholomorphic involution on the periods of $CY_3(3,243)$, given its action on its cohomology, using a canonical (co)homology basis. Section **6** has the conclusions and discussion. There is one appendix on the evaluation of the monodromy matrix.

2 Landau-Ginsburg PF equation for type IIA on $CY_3(3,243)$

By following the alternative formulation of Hori and Vafa [1] for deriving the Picard-Fuchs equation for a definition of period integral in the mirror Landau-Ginsburg model, we obtain solutions valid in the large *and* small complex structure limits, and get the ln terms as naturally as the analytic terms (i.e. without using parametric differentiation of infinite series). We also study in detail, the monodromy matrix in the large and small complex structure limits.

Consider the Calabi-Yau 3-fold given as a degree-24 Fermat hypersurface in the weighted projective space $\mathbf{WCP}^4[1,1,2,8,12]$:

$$P = z_1^{24} + z_2^{24} + z_3^{12} + z_4^3 + z_5^2 = 0. \tag{1}$$

It has a $\mathbf{Z}_2$-singularity curve and a $\mathbf{Z}_4$-singularity point. $\mathbf{Z}_2$ and $\mathbf{Z}_4$ singularity resolution ↔ The two new chiral superfields needed to be introduced as a consequence of singularity resolution, correspond to the two $\mathbf{CP}^1$'s that required to be introduced in blowing up the singularities. One then has to consider three instead of a single C^* action, and the $CY_3(3,243)$ [2] can be expressed as a suitable holomorphic quotient corresponding to a smooth toric variety. To be more specific, one considers the resolved Calabi-Yau $CY_3(3,243)$ as the holomorphic quotient: $\frac{C^7-F}{(C^*)^3}|_{\text{hyp constraint}}$, where the diagonal $(C^*)^3$ actions on the seven coordinates of C^7 are given by:

$$x^j \sim \lambda^{iQ^a_j} x^j, \ \text{no sum over } j; \ a = 1,2,3, \tag{2}$$

where the three sets of charges $\{Q^{a=1,2,3}_{i=(0,),1,\ldots,7}\}$ (the "0" being for the extra chiral superfield with $Q^0_i = -\sum_{i=1}^7 Q^a_i[7]$) are give by the following:

$$\begin{array}{c|cccccccc}
 & \mathcal{X}_0 & \mathcal{X}_1 & \mathcal{X}_2 & \mathcal{X}_3 & \mathcal{X}_4 & \mathcal{X}_5 & \mathcal{X}_6 & \mathcal{X}_7 \\
\hline
Q^{(1)}_i : & 0 & 1 & 1 & -2 & 0 & 0 & 0 & 0 \\
Q^{(2)}_i : & 0 & 0 & 0 & 1 & 1 & 0 & 0 & -2 \\
Q^{(3)}_i : & -6 & 0 & 0 & 0 & 0 & 2 & 3 & 1
\end{array} \tag{3}$$

where on noting:

$$Q^{(1)} + 2Q^{(2)} + 4Q^{(3)} = \ -24 \ \ 1 \ \ 1 \ \ 0 \ \ 2 \ \ 8 \ \ 12 \ \ 0 \ , \tag{4}$$

---

[2] The $CY_3(3,243)$ considered in this paper will be an elliptic fibration over the Hirzebruch surface F_2.

one identifies $\mathcal{X}_{3,7}$ as the two extra chiral superfields introduced as a consequence of singularity resolution.

The Landau-Ginsburg Period for the resolved $CY_3(3,243)$, as per the prescription of Hori and Vafa, is given by:

$$\begin{aligned}
\Pi(t_1,t_2,t_3) &= \int \prod_{i=0}^{7} dY_i \prod_{a=1}^{3} dF^{(a)} \sum_{a=1}^{3} d_{1a} F^{(a)} e^{-\sum_{a=1}^{3} F^{(a)}(\sum_{i=1}^{7} Q_i^{(a)} Y_i - Q_0^{(a)} Y_0 - t_{(a)}) - \sum_{i=0}^{7} e^{-Y_i}} \\
&= \sum_{a=1}^{3} d_{1a} \frac{\partial}{\partial t_{(a)}} \int \prod_{i=0}^{7} dY_i \prod_{a=1}^{3} \delta(\sum_{i=1}^{7} Q_i^{(a)} Y_i - Q_0^{(a)} Y_0 - t_{(a)}) e^{-\sum_{i=0}^{7} e^{-Y_i}} \\
&\equiv \sum_{a=1}^{3} d_{1a} \frac{\partial}{\partial t_{(a)}} \tilde{\Pi}(t_{1,2,3}); \qquad (5)
\end{aligned}$$

$d_{\alpha a} \equiv$ charge matrix, α indexes the number of hypersurfaces and a indexes the number of $U(1)$'s. For $CY_3(3,243)$, $\alpha = 1$, $a = 1,2,3$ with $d_{11} = d_{12} = 0, d_{13} = 6$.

Consider:

$$\tilde{\Pi}(t_{1,2,3}, \{\mu_i\}) \equiv \int \prod_{i=0}^{7} dY_i \prod_{a=1}^{3} \delta(\sum_{i=1}^{7} Q_i^{(a)} Y_i - Q_0^{(a)} Y_0 - t_{(a)}) e^{-\sum_{i=0}^{7} \mu_i e^{-Y_i}}. \quad (6)$$

One can show that:

$$\tilde{\Pi}(t_{1,2,3}, \{\mu_i\}) = \tilde{\Pi}(t'_{1,2,3}, \{\mu_i = 1\}), \quad (7)$$

where

$$t'_1 \equiv t_1 + ln(\mu_3^2/\mu_1\mu_2),\ t'_2 = t_2 + ln(\mu_7^2/\mu_3\mu_4),\ t'_3 = t_3 + ln(\mu_0^6/\mu_7\mu_5^2\mu_6^3). \quad (8)$$

Eliminating $Y_{0,3,7}$ gives a order-24 Picard-Fuchs equation:

$$\frac{\partial^{24}}{\partial\mu_1\partial\mu_2\partial\mu_4^2\partial\mu_5^8\partial\mu_6^{12}} \tilde{\Pi}(t_{1,3,4}) = e^{-t_1+2t_2+4t_3} \frac{\partial^{24}}{\partial\mu_0^{24}} \tilde{\Pi}(t_{1,2,3}), \quad (9)$$

which is the same as the PF equation for the unresolved hypersurface away from the orbifold singularities. **This overcounts the number of solutions**.

The right number of solutions must be $2h^{2,1}(\text{Mirror}) + 2 = 2.3 + 2 = 8$. To get this number, one notes that by adding the three constraints:

$$Y_1 + Y_2 - 2Y_3 = t_1;\ Y_3 + Y_4 - 2Y_7 = t_2;\ -6Y_0 + 2Y_5 + 3Y_6 + Y_7 = t_3, \quad (10)$$

one gets:

$$-6Y_0 - Y_3 - Y_7 + Y_1 + Y_2 + Y_4 + 2Y_5 + 3Y_6 = t_1 + t_2 + t_3, \quad (11)$$

which allows one to write the following order-8 PF equation:

$$\frac{\partial^8}{\partial\mu_1\partial\mu_2\partial\mu_4\partial\mu_5^2\partial\mu_6^3}\tilde{\Pi}(t_{1,2,3}) = e^{-(t_1+t_2+t_3)}\frac{\partial^8}{\partial\mu_0^6\partial\mu_3\partial\mu_7}\tilde{\Pi}(t_{1,2,3}). \tag{12}$$

If $\Theta_i \equiv \frac{\partial}{\partial t'_i}$, then one gets:

$$\Big[\Theta_1^2\Theta_2\prod_{l=2}^{3}\prod_{k=0}^{l-1}(-l\Theta_3 - k) - e^{-(t'_1+t'_2+t'_3)}(2\Theta_2 - \Theta_3)(2\Theta_1 - \Theta_2)\prod_{j=0}^{5}(6\Theta_3 - j)\Big]\tilde{\Pi} = 0$$

with $z \equiv e^{-(t'_1+t'_2+t'_3)}$; $z\dfrac{d}{dz} \equiv \Delta_z$, and rescaling :

$$\Big[\Delta_z^4(\Delta_z - \frac{1}{2})\Delta_z(\Delta_z - \frac{1}{3})(\Delta_z - \frac{2}{3}) + z\prod_{j=0}^{5}(\Delta_z + \frac{j}{6})\Delta_z^2\Big]\tilde{\Pi} = 0. \tag{13}$$

One solution to the above equation is: ${}_8F_7\left(\begin{array}{cccccccc} 0 & \frac{1}{6} & \frac{2}{6} & \frac{3}{6} & \frac{4}{6} & \frac{5}{6} & 0 & 0 \\ 1 & 1 & 1 & \frac{1}{2} & 1 & \frac{2}{3} & \frac{1}{3} & \end{array}\right)(-z)$

[For $e^{-t'} \equiv z$, $t' \equiv t'_1 + 2t'_2 + 4t'_3$ and suitable rescaling of z, the relevant order-24 PF equation **for the unresolved hypersurface** is:

$$\Delta_z^2\Delta_z(\Delta_z - \frac{1}{2})\prod_{j=1}^{8}(\Delta_z - \frac{j-1}{8})\prod_{j=1}^{12}(\Delta_z - \frac{j-1}{12})\tilde{\Pi} = z\prod_{j=1}^{24}(\Delta_z + \frac{j-1}{24})\tilde{\Pi}. \tag{14}$$

One solution can be written in terms of the following generalized hypergeometric function

$${}_{24}F_{23}\left(\begin{array}{cccccccccc} 0 & \frac{1}{24} & \frac{2}{24} & \frac{3}{24} & & \frac{4}{24} & & \frac{5}{24} & .... & \frac{23}{24} \\ 1 & 1 & \frac{1}{2} & \frac{5}{8} & ... & - & \frac{2}{8} & \frac{1}{12} & ... & -\frac{10}{12} \end{array}\right). \tag{15}$$

]

From the above solution, Meijer basis obtained using properties of ${}_pF_q$ and the Meijer function I:

$${}_pF_q\left(\begin{array}{ccccc} \alpha_1 & \alpha_2 & \alpha_3 & .... & \alpha_p \\ \beta_1 & \beta_2 & \beta_3 & .... & \beta_q \end{array}\right)(z) = \frac{\prod_{i=1}^{p}\Gamma(\beta_i)}{\prod_{j=1}^{q}\Gamma(\alpha_j)} I\left(\begin{array}{c|c} 0 & \alpha_1...\alpha_p \\ \hline . & \beta_1...\beta_q \end{array}\right)(-z) \text{ where}$$

$$I\left(\begin{array}{c|c} a_1...a_A & b_1...b_B \\ \hline c_1...c_C & d_1...d_D \end{array}\right)(z), I\left(\begin{array}{c|c} a_1...(1-d_l)...a_A & b_1...b_B \\ \hline c_1...c_C & d_1...\hat{d}_l...d_D \end{array}\right)(-z)$$

$$I\left(\begin{array}{c|c} a_1...a_A & b_1..\hat{b}_j...b_B \\ \hline c_1..(1-b_j)..c_C & d_1...d_D \end{array}\right)(-z) \tag{16}$$

satisfy the same equation.

Now, $z \equiv e^{-(t'_1+t'_2+t'_3)} = e^{-(t_1+t_2+t_3)}\frac{\mu_1\mu_2\mu_4\mu_5^2\mu_6^3}{\mu_0^6\mu_3\mu_7}$. Hence, one can solve for large ($\equiv |z| << 1$) and small complex structure ($\equiv |z| >> 1$) limits, as well as large-size-Calabi-Yau limit($\equiv t_i \rightarrow \infty \Leftrightarrow |z| << 1$) on the mirror Landau-Ginsburg side with equal ease using Mellin-Barnes integral represention for the Meijer's function I, as in [8] and in (17) below.

Now, to get an infinite series expansion in z for $|z| < 1$ as well as $|z| > 1$, one uses the following

$$I\left(\begin{array}{c|c} a_1...a_A & b_1...b_B \\ \hline c_1...c_C & d_1...d_D \end{array}\right)(z) = \frac{1}{2\pi i}\int_\gamma ds \frac{\prod_{i=1}^A \Gamma(a_i - s)\prod_{j=1}^B \Gamma(b_j + s)}{\prod_{k=1}^C \Gamma(c_k - s)\prod_{l=1}^D \Gamma(d_l + s)} z^s, \quad (17)$$

where the contour γ lies to the right of:$s + b_j = -m \in \mathbf{Z}^- \cup \{0\}$ and to the left of: $a_i - s = -m \in \mathbf{Z}^- \cup \{0\}$.



Figure 1: The contour γ for I: The poles $s = a_i + m$ are denoted by # and the poles $s = -m - b_j$ are denoted by x

This, $|z| << 1$ and $|z| >> 1$ can be dealt with equal ease by suitable deformations of the contour γ (see Fig. 1) to γ' and γ'' respectively (See Fig. 2). Additionally, instead of performing parametric differentiation of infinite series to get the ln-terms, one get the same (for the large complex structure limit: $|z| < 1$) by evaluation of the residue at $s = 0$ in the Mellin-Barnes contour integral in (17) as is done explicitly to evaluate the eight integrals



Figure 2: The deformed contour γ' valid for $|z| < 1$, and γ'' valid for $|z| >> 1$

in (19). [3]

The guiding principle is that of the eight solutions to $\tilde{\Pi}$, one should generate solutions in which one gets $(lnz)^P$, $P = 1, ..., 4$ so that one gets $(lnz)^{P-1}$ for Π, and one can then identify terms independent of lnz with Z^0, three (lnz) terms with $Z^{1,2,3}$, three $(lnz)^{P\leq 2}$ terms with $F_{1,2,3} \equiv \frac{\partial F}{\partial Z^{1,2,3}}$, and finally $(lnz)^{P\leq 3}$ term with $F_0 \equiv \frac{\partial F}{\partial Z^0}$.

---

[3] From [2], one gets, for the case of large-base Calabi-Yau away from the orbifold singularities:

$$
\begin{aligned}
& I\left(\begin{array}{c|c} 0 & 0\frac{1}{24}\cdots\frac{23}{24} \\ \hline . & 11\frac{1}{2}\frac{5}{8}\cdots-\frac{2}{8}\frac{1}{12}\cdots-\frac{10}{12} \end{array}\right)(-z) = \Theta(1-|z|)\sum_{m=1}^{\infty}\frac{\prod_{j=1}^{23}\Gamma(m+\frac{j}{24})z^m(-)^{m+1}}{\Gamma(m+\frac{1}{2})\prod_{k=-2}^{5}\Gamma(m+\frac{k}{8})\prod_{l=-10}^{1}\Gamma(m+\frac{l}{10})m^2m!(m-1)!} \\
& \times\Big[\sum_{j=1}^{23}\Psi(m+\frac{j}{24})+ln(-z)-1-\Psi(\frac{1}{2}+m)-\sum_{k=-2}^{5}\Psi(m+\frac{k}{8})-\sum_{l=10}^{2}\Psi(\frac{l}{12}+m)-\frac{1}{m}-\sum_{j=1}^{m}\frac{1}{j}-\Psi(m+1)\Big] \\
& +\Theta(|z|-1)\sum_{m=0}^{\infty}\sum_{j=1}^{23}\frac{-\Gamma(\frac{j}{24}+m)(-)^m\prod_{j\neq k=1}^{23}\Gamma(\frac{k-j}{24}-m)}{m!(m+\frac{j}{24})\Gamma(-m-\frac{j}{24}+1)\prod_{k=-2}^{8}\Gamma(\frac{k}{8}-m-\frac{j}{24})\prod_{l=10}^{1}\Gamma(\frac{l}{12}-m-\frac{j}{24})}(-z)^{-m-\frac{j}{24}} \\
& =\Theta(1-|z|)\sum_{m=1}^{\infty}(2\pi)^{\frac{7}{2}}\frac{24^{\frac{1}{2}-24m}}{8^{\frac{1}{2}-8[m-\frac{1}{4}]}.10^{\frac{1}{2}-10[m-1]}}\frac{\Gamma(24m)z^m(-)^{m+1}}{\Gamma(m+\frac{1}{2})\Gamma(m-\frac{1}{10})\Gamma(10m)m^2(m-1)!} \\
& \times\left[-\Psi(m)+24\Psi(24m)+ln\left(\frac{12^{12}.8^8}{24^{24}}\right)+ln(-z)-1-\Psi(\frac{1}{2}+m)-8\Psi(8[m-\frac{1}{4}])-12\Psi(12[m-\frac{5}{6}])\right. \\
& \left.-\frac{1}{m}-\sum_{j=1}^{m}\frac{1}{j}-\Psi(m+1)\right] \\
& -\Theta(|z|-1) \\
& \times\sum_{m=0}^{\infty}\sum_{j=1}^{23}\frac{\Gamma(\frac{j}{24}+m)\Gamma(m+\frac{j}{24}+\frac{1}{4})\Gamma(m+\frac{j}{24}+\frac{5}{6})(-)^m(-z)^{-m-\frac{j}{24}}\prod_{j\neq k}^{23}\Gamma(\frac{k-j}{24}-m)}{m!(m+\frac{j}{24})\Gamma(-m-\frac{j}{24}+1).(2\pi)^9.8^{\frac{1}{2}-8[m+\frac{j}{24}+\frac{1}{4}]}.12^{\frac{1}{2}-12[m+\frac{j}{24}+\frac{5}{6}]}\Gamma(8[m+\frac{j}{24}+\frac{1}{4}])\Gamma(12[m+\frac{j}{24}+\frac{5}{6}])}
\end{aligned}
\tag{18}
$$

One (non-unique) choice of solutions for $\tilde{\Pi}(z)$ is given below:

$$
z\frac{d}{dz}\left[\begin{array}{c}
I\left(\begin{array}{c|c} 0\ 0 & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0\ 0 \\ \hline . & 1\ 1\ \frac{1}{2}\ 1\ \frac{2}{3}\ \frac{1}{3} \end{array}\right)(-z) \\
I\left(\begin{array}{c|c} 0\ 0 \frac{1}{2}\ \frac{1}{3}\ \frac{2}{3} & \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6} \\ \hline 1\ 1\ 1 & 1\ 1\ 1 \end{array}\right)(-z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{1}{2} & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ 1\ \frac{2}{3}\ \frac{1}{3} \end{array}\right)(-z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{1}{3} & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ \frac{1}{2}\ 1\ \frac{1}{3} \end{array}\right)(z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{2}{3} & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ \frac{1}{2}\ 1\ \frac{2}{3} \end{array}\right)(-z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{1}{2} & \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ 1\ \frac{2}{3}\ \frac{1}{3} \end{array}\right)(z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{1}{3} & \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ \frac{1}{2}\ \frac{2}{3}\ \frac{1}{3} \end{array}\right)(z) \\
I\left(\begin{array}{c|c} 0\ 0\ \frac{2}{3} & \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ \frac{1}{2}\ \frac{2}{3} \end{array}\right)(z)
\end{array}\right] \sim \left[\begin{array}{c} F_0 \\ Z^0 \\ F_1 \\ F_2 \\ F_3 \\ Z^1 \\ Z^2 \\ Z^3 \end{array}\right] \qquad (19)
$$

(a)

$$I\left(\begin{array}{c|c} 0\ 0 & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0\ 0 \\ \hline . & 1\ 1\ \frac{1}{2}\ 1\ \frac{2}{3}\ \frac{1}{3} \end{array}\right)(-z) = \frac{1}{2\pi i}\int_\gamma ds \frac{[\Gamma(-s)]^2[\Gamma(s)]^3\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{[\Gamma(s+1)]^3\Gamma(\frac{1}{2}+s)\Gamma(\frac{2}{3}+s)\Gamma(\frac{1}{3}+s)}(-z)^s$$

$$= \theta(1-|z|)\Bigg[\frac{2(2\pi)^{\frac{3}{2}}}{\sqrt{\pi}}\Bigg[\Bigg(ln\Big(\frac{2^2 3^3}{6^6}\Big)+ln(-z)\Bigg)^4+\frac{65}{3}\Bigg(ln\Big(\frac{2^2 3^3}{6^6}\Big)+ln(-z)\Bigg)^2$$

$$+\frac{169\pi^4}{3}-1440\zeta(3)\Bigg(ln\Big(\frac{2^2 3^3}{6^6}\Big)+ln(-z)\Bigg)+402\zeta(4)\Bigg]+\sum_{m=1}^\infty\frac{(2\pi)^{\frac{3}{2}}2^{\frac{1}{2}-6m}\Gamma(6m)}{m^3(m!)^2 3^{3m}\Gamma(\frac{1}{2}+m)\Gamma(3m)}$$

$$\times\Bigg[2\gamma+6\Psi(6m)-2\Psi(2m)-3\Psi(m)-3m^2+ln\Big(\frac{2^2 3^3}{6^6}\Big)\Bigg](-z)^m\Bigg]$$

$$-\theta(|z|-1)\Bigg[\sum_{m=0}^\infty\frac{(-)^m(\Gamma(m+\frac{1}{6})^2\Gamma(-m+\frac{2}{3})}{(m+\frac{1}{6})^3 m!}(-z)^{-m-\frac{1}{6}}+$$

$$\sum_{m=0}^\infty\frac{(-)^m(\Gamma(m+\frac{5}{6})^2\Gamma(-m-\frac{2}{3})}{(m+\frac{5}{6})^3 m!}(-z)^{-m-\frac{5}{6}}\Bigg] \qquad (20)$$

(b)

$$I\left(\begin{array}{c|c} 0\ 0\frac{1}{2}\ \frac{1}{3}\ \frac{2}{3} & \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6} \\ \hline 1\ 1\ 1 & 1\ 1\ 1 \end{array}\right)(-z)$$

$$= \frac{1}{2\pi i}\int_\gamma ds\frac{[\Gamma(-s)]^2\Gamma(\frac{1}{2}-s)\Gamma(\frac{1}{3}-s)\Gamma(\frac{2}{3}-s)\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{[\Gamma(1-s)]^3[\Gamma(1+s)]^2}(-z)^s$$

$$= \Theta(1-|z|)\Bigg[\frac{\sqrt{\pi}.2\pi.(2\pi)^{\frac{5}{2}}}{\sqrt{18}}\Bigg(ln[\frac{2^2 3^3}{6^6}]+ln(-z)\Bigg)$$

$$+\sum_{m=0}^\infty\frac{(-)^{m+1}(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6[m+\frac{1}{2}]}}{m!(m+\frac{1}{2})^2}\frac{\Gamma(\frac{1}{3}-[\frac{1}{2}+m])\Gamma(\frac{2}{3}-[\frac{1}{2}+m])\Gamma(6[m+\frac{1}{2}])}{[\Gamma(\frac{3}{2}-m)]^2\Gamma(\frac{1}{2}-m)\Gamma(m+\frac{1}{2})}(-z)^{m+\frac{1}{2}}$$

$$+\sum_{m=0}^\infty\frac{(-)^{m+1}(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6[m+\frac{1}{3}]}}{m!(m+\frac{1}{3})^2}\frac{\Gamma(\frac{1}{2}-[\frac{1}{3}+m])\Gamma(\frac{2}{3}-[\frac{1}{3}+m])\Gamma(6[m+\frac{1}{3}])}{[\Gamma(\frac{2}{3}-m)][\Gamma(\frac{2}{3}+m)]^2\Gamma(m+\frac{1}{3})}(-z)^{m+\frac{1}{3}}$$

$$+\sum_{m=0}^\infty\frac{(-)^{m+1}(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6[m+\frac{2}{3}]}}{m!(m+\frac{2}{3})^2}\frac{\Gamma(\frac{1}{2}-[\frac{2}{3}+m])\Gamma(\frac{1}{3}-[\frac{2}{3}+m])\Gamma(6[m+\frac{2}{3}])}{[\Gamma(\frac{1}{3}-m)][\Gamma(\frac{5}{3}+m)]^2\Gamma(m+\frac{2}{3})}(-z)^{m+\frac{2}{3}}\Bigg]$$

$$+\Theta(|z|-1)$$

$$\times\Bigg[\sum_{m=0}^\infty\sum_{j=1}^5\frac{(-)^{m+1}}{m!(m+\frac{j}{6})^2}\frac{\Gamma(\frac{1}{2}+m+\frac{j}{6}\Gamma(\frac{1}{3}+m+\frac{j}{6})\Gamma(\frac{2}{3}+m+\frac{j}{6})\prod_{k\neq j}\Gamma(-m+\frac{k-j}{6})}{\Gamma(1+m+\frac{j}{6})[\Gamma(1-m-\frac{j}{6})]^2}(-z)^{-m-\frac{j}{6}}\Bigg]$$

$$\tag{21}$$

(c)

$$I\left(\begin{array}{c|c} 0\ 0\tfrac{1}{2} & \tfrac{1}{6}\ \tfrac{2}{6}\ \tfrac{3}{6}\ \tfrac{4}{6}\ \tfrac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ 1\ \tfrac{2}{3}\ \tfrac{1}{3} \end{array}\right)(z)$$
$$= \frac{1}{2\pi i}\int_\gamma ds\frac{[\Gamma(-s)]^2\Gamma(\frac{1}{2}-s)\Gamma(s)\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{[\Gamma(1-s)]^2[\Gamma(1+s)]^3\Gamma(\frac{2}{3}+s)\Gamma(\frac{1}{3}+s)}(z)^s$$
$$\Theta(1-|z|)\Bigg[2\pi^2\Big[(lnz+ln\Big(\frac{2^2.3^3}{6^6}\Big)^2+\frac{14\pi^2}{3}\Big]$$
$$+\sum_{m=0}^{\infty}\frac{(-)^{m+1}z^{m+\frac{1}{2}}}{m!}\frac{(2\pi)^{\frac{5}{2}}.6^{\frac{1}{2}-6[m+\frac{1}{2}]}\Gamma(6[m+\frac{1}{2}])}{(m+\frac{1}{2})^3[\Gamma(\frac{3}{2}+m)]^2.2\pi.3^{\frac{1}{2}-3[m+\frac{1}{2}]}\Gamma(3[m+\frac{1}{2}])}[\Gamma(\frac{3}{2}+m)]^2\Bigg]$$
$$-\Theta(|z|-1)\Bigg[\sum_{m=0}^{\infty}\frac{(-)^{m+1}.z^{-m-\frac{j}{6}}}{m!}\frac{\Gamma(\frac{1}{2}+m+\frac{j}{6})\prod_{k\neq j}\Gamma(\frac{k-j}{6}-m)}{(m+\frac{j}{6})^3[\Gamma(1-m-\frac{j}{6})]^2\Gamma(\frac{2}{3}-m-\frac{j}{6})\Gamma(\frac{1}{3}-m-\frac{j}{6})}\Bigg] \tag{22}$$

(d)

$$I\left(\begin{array}{c|c} 0\ 0\tfrac{1}{3} & \tfrac{1}{6}\ \tfrac{2}{6}\ \tfrac{3}{6}\ \tfrac{4}{6}\ \tfrac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ \tfrac{1}{2}\ 1\ \tfrac{1}{3} \end{array}\right)(z)$$
$$= \frac{1}{2\pi i}\int_\gamma ds\frac{[\Gamma(-s)]^2\Gamma(\frac{1}{3}-s)\Gamma(s)\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{[\Gamma(1-s)]^2[\Gamma(1+s)]^3\Gamma(\frac{1}{2}+s)\Gamma(\frac{1}{3}+s)}(z)^s$$
$$=\Theta(1-|z|)\Bigg[\sqrt{\frac{16}{3}}.\pi^2\Big[\Big(ln\Big[\frac{2^2.3^3}{6^6}\Big]+lnz+\frac{\pi}{\sqrt{3}}\Big)^2+5\pi^2\Big]$$
$$+\sum_{m=0}^{\infty}\frac{(-)^{m+1}}{m!}\frac{(2\pi)^{\frac{5}{2}}.6^{\frac{1}{2}-6[\frac{1}{3}+m]}\Gamma(6[\frac{1}{3}+m])}{(m+\frac{1}{3})^3\Gamma(\frac{1}{3}+m)[\Gamma(\frac{4}{3}+m)]^2\Gamma(\frac{5}{6}+m)\Gamma(\frac{2}{3}+m)}\Bigg]$$
$$-\Theta(|z|-1)\sum_{m=0}^{\infty}\sum_{j=1}^{5}\frac{(-)^{m+1}}{m!}\frac{\Gamma(\frac{1}{3}+m+\frac{j}{6})\prod_{k\neq j}^5\Gamma(\frac{k-j}{6}-m)}{(\frac{j}{6}+m)^3[\Gamma(1-\frac{j}{6}-m)]^2\Gamma(\frac{1}{2}-\frac{j}{6}-m)\Gamma(\frac{1}{3}-\frac{j}{6}-m)}z^{-\frac{j}{6}-m} \tag{23}$$

(e)

$$
\begin{aligned}
&I\left(\begin{array}{c|c} 0\ 0\tfrac{2}{3} & \tfrac{1}{6}\ \tfrac{2}{6}\ \tfrac{3}{6}\ \tfrac{4}{6}\ \tfrac{5}{6}\ 0 \\ \hline 1\ 1 & 1\ 1\ \tfrac{1}{2}\ 1\ \tfrac{2}{3} \end{array}\right)(z) \\
&= \frac{1}{2\pi i}\int_\gamma ds \frac{[\Gamma(-s)]^2\Gamma(\frac{2}{3}-s)\Gamma(s)\prod_{j=1}^{5}\Gamma(s+\frac{j}{6})}{[\Gamma(1-s)]^2[\Gamma(1+s)]^3\Gamma(\frac{1}{2}+s)\Gamma(\frac{2}{3}+s)}(z)^s \\
&= \Theta(1-|z|)\Bigg[\sqrt{\frac{16}{3}}.\pi^2\Bigg[\Bigg(ln\Big[\frac{2^2.3^3}{6^6}\Big]+lnz-\frac{\pi}{\sqrt{3}}\Bigg)^2+5\pi^2\Bigg] \\
&+\sum_{m=0}^{\infty}\frac{(-)^{m+1}}{m!}\frac{(2\pi)^{\frac{5}{2}}.6^{\frac{1}{2}-6[\frac{2}{3}+m]}\Gamma(6[\frac{2}{3}+m])}{(m+\frac{1}{3})^3\Gamma(\frac{2}{3}+m)[\Gamma(\frac{4}{3}+m)]^2\Gamma(\frac{7}{6}+m)\Gamma(\frac{4}{3}+m)}\Bigg] \\
&-\Theta(|z|-1)\sum_{m=0}^{\infty}\sum_{j=1}^{5}\frac{(-)^{m+1}}{m!}\frac{\Gamma(\frac{2}{3}+m+\frac{j}{6})\prod_{k\neq j}^{5}\Gamma(\frac{k-j}{6}-m)}{(\frac{j}{6}+m)^3[\Gamma(1-\frac{j}{6}-m)]^2\Gamma(\frac{1}{2}-\frac{j}{6}-m)\Gamma(\frac{2}{3}-\frac{j}{6}-m)}z^{-\frac{j}{6}-m}
\end{aligned}
\tag{24}
$$

(f)

$$
\begin{aligned}
&I\left(\begin{array}{c|c} 0\ 0\ \tfrac{1}{2} & 0\ \tfrac{1}{6}\ \tfrac{2}{6}\ \tfrac{3}{6}\ \tfrac{4}{6}\ \tfrac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ 1\ \tfrac{2}{3}\ \tfrac{1}{3} \end{array}\right)(-z) \\
&= \frac{1}{2\pi i}\int_\gamma \frac{[\Gamma(-s)]^2[\Gamma(s)]^2\Gamma(\frac{1}{2}-s)\prod_{j=1}^{5}\Gamma(s+\frac{j}{6})}{\Gamma(1-s)[\Gamma(1+s)]^3\Gamma(s+\frac{2}{3})\Gamma(s+\frac{1}{3})}(-z)^s \\
&= \Theta(1-|z|)\Bigg[2\pi^2\Bigg(\Bigg(ln(-z)+ln\Big[\frac{2^3.3^3}{6^6}\Big]\Bigg)^3+15\pi^2\Bigg(ln(-z)+ln\Big[\frac{2^2.3^3}{6^6}\Big]\Bigg)-356\zeta(3)\Bigg) \\
&+\sum_{m=1}^{\infty}\frac{(-)^m(2\pi)^{\frac{3}{2}}}{(m!)^2m^3}\frac{6^{\frac{1}{2}-6m}}{3^{\frac{1}{2}-3m}}\frac{\Gamma(6m)}{\Gamma(3m)}\Gamma(\frac{1}{2}-m)(-z)^m \\
&+\sum_{m=1}^{\infty}\frac{(-)^m}{(m!)^2(m+\frac{1}{2})^3}.\frac{(2\pi)^{\frac{3}{2}}.6^{\frac{1}{2}-6[m+\frac{1}{2}]}}{3^{\frac{1}{2}-3[m+\frac{1}{2}]}}\frac{\Gamma(6[m+\frac{1}{2}])}{\Gamma(3[m+\frac{1}{2}])}(-z)^{m+\frac{1}{2}}\Bigg] \\
&+\Theta(|z|-1)\sum_{m=0}^{\infty}\sum_{j=1}^{5}\frac{(-)^m}{2\pi 3^{\frac{1}{2}+3[\frac{j}{6}+m)]}m!(\frac{j}{6}+m)^3}\frac{\Gamma(\frac{j}{6}+m)}{\Gamma(1-\frac{j}{6}-m)}.\Gamma(\frac{1}{2}+\frac{j}{6}+m).\frac{\Gamma(-[\frac{j}{6}+m])}{\Gamma(-3[\frac{j}{6}+m])} \\
&\times\prod_{k\neq j}\Gamma(\frac{k-j}{6}-m).z^{-\frac{j}{6}-m}
\end{aligned}
\tag{25}
$$

(g)

$$
\begin{aligned}
&I\left(\begin{array}{c|c} 0\ 0\ \frac{1}{3} & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ \frac{1}{2}\ 1\ \frac{1}{3} \end{array}\right)(-z) \\
&= \frac{1}{2\pi i}\int_\gamma \frac{[\Gamma(-s)]^2[\Gamma(s)]^2\Gamma(\frac{1}{3}-s)\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{\Gamma(1-s)[\Gamma(1+s)]^3\Gamma(\frac{1}{2}+s)\Gamma(\frac{1}{3}+s)}(-z)^s \\
&= \Theta(1-|z|)\Bigg[\pi^2\sqrt{\frac{16}{3}}\Bigg(\Bigg[ln\Bigg(\frac{2^2.3^3}{6^6}\Bigg)+ln(-z)+\frac{\pi}{\sqrt{3}}\Bigg)^3 \\
&+16\pi^2\Bigg[ln\Bigg(\frac{2^2.3^3}{6^6}\Bigg)+ln(-z)+\frac{\pi}{\sqrt{3}}\Bigg]-360\zeta(3)-\frac{8\pi^3}{3^{\frac{3}{2}}}\Bigg) \\
&+\sum_{m=1}^\infty \frac{(-)^{m+1}}{(m!)^2m^3}\frac{\Gamma(\frac{1}{3}-m)}{\Gamma(\frac{1}{3}+m)}(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6m}\frac{\Gamma(6m)}{\Gamma(m)\Gamma(\frac{1}{2}+m)}(-z)^m \\
&+\sum_{m=0}^\infty \frac{(-)^{m+1}}{m!(\frac{1}{3}+m)^4}\frac{\Gamma(\frac{2}{3}-m)}{\Gamma(\frac{4}{3}+m)}\frac{(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6[\frac{1}{3}+m]}\Gamma(6[\frac{1}{3}+m])}{\Gamma(\frac{1}{3}+m)}\frac{(-z)^{m+\frac{1}{3}}}{\Gamma(\frac{5}{6}+m)\Gamma(\frac{2}{3}+m)}\Bigg] \\
&+\Theta(|z|-1)\sum_{m=0}^\infty\sum_{j=1}^5 \frac{(-)^{m+1}}{m!(\frac{j}{6}+m)^4}\frac{\Gamma(\frac{1}{3}+\frac{j}{6}+m)\Gamma(1+\frac{j}{6}+m)}{\Gamma(1-\frac{j}{6}-m)\Gamma(\frac{1}{2}-\frac{j}{6}-m)}\frac{\prod_{k\neq j}\Gamma(\frac{k-j}{6}-m)}{\Gamma(\frac{1}{3}-\frac{j}{6}-m)}(-z)^{-\frac{j}{6}-m}
\end{aligned}
\tag{26}
$$

(h)

$$
\begin{aligned}
&I\left(\begin{array}{c|c} 0\ 0\ \frac{2}{3} & 0\ \frac{1}{6}\ \frac{2}{6}\ \frac{3}{6}\ \frac{4}{6}\ \frac{5}{6}\ 0 \\ \hline 1 & 1\ 1\ \frac{1}{2}\ 1\ \frac{2}{3} \end{array}\right)(-z) \\
&= \frac{1}{2\pi i}\int_\gamma \frac{[\Gamma(-s)]^2[\Gamma(s)]^2\Gamma(\frac{2}{3}-s)\prod_{j=1}^5\Gamma(s+\frac{j}{6})}{\Gamma(1-s)[\Gamma(1+s)]^3\Gamma(\frac{1}{2}+s)\Gamma(\frac{2}{3}+s)}(-z)^s \\
&= \Theta(1-|z|)\Bigg[\pi^2\sqrt{\frac{16}{3}}\Bigg(\Bigg[ln\Bigg(\frac{2^2.3^3}{6^6}\Bigg)+ln(-z)-\frac{\pi}{\sqrt{3}}\Bigg)^3 \\
&+16\pi^2\Bigg[ln\Bigg(\frac{2^2.3^3}{6^6}\Bigg)+ln(-z)+\frac{\pi}{\sqrt{3}}\Bigg]-360\zeta(3)+\frac{8\pi^3}{3^{\frac{3}{2}}}\Bigg) \\
&+\sum_{m=1}^\infty \frac{(-)^{m+1}}{(m!)^2m^3}\frac{\Gamma(\frac{2}{3}-m)}{\Gamma(\frac{1}{3}+m)}(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6m}\frac{\Gamma(6m)}{\Gamma(m)\Gamma(\frac{1}{2}+m)}(-z)^m \\
&+\sum_{m=0}^\infty \frac{(-)^{m+1}}{m!(\frac{2}{3}+m)^4}\frac{\Gamma(\frac{1}{3}-m)}{\Gamma(\frac{5}{3}+m)}\frac{(2\pi)^{\frac{5}{2}}6^{\frac{1}{2}-6[\frac{2}{3}+m]}\Gamma(6[\frac{2}{3}+m])}{\Gamma(\frac{2}{3}+m)}\frac{(-z)^{m+\frac{2}{3}}}{\Gamma(\frac{7}{6}+m)\Gamma(\frac{4}{3}+m)}\Bigg]
\end{aligned}
$$

$$+\Theta(|z|-1)\sum_{m=0}^{\infty}\sum_{j=1}^{5}\frac{(-)^{m+1}}{m!(\frac{j}{6}+m)^4}\frac{\Gamma(\frac{2}{3}+\frac{j}{6}+m)\Gamma(1+\frac{j}{6}+m)}{\Gamma(1-\frac{j}{6}-m)\Gamma(\frac{1}{2}-\frac{j}{6}-m)}\frac{\prod_{k\neq j}\Gamma(\frac{k-j}{6}-m)}{\Gamma(\frac{2}{3}-\frac{j}{6}-m)}(-z)^{-\frac{j}{6}-m} \tag{27}$$

[4]

The connection between (19) that *effectively depends only on one complex structure parameter* $z = e^{-(t_1+t_2+t_3)}\frac{1}{z_1 z_2 z_3}$, and the solutions given in the literature [9]of the form:

$$\partial^{s_m}_{\rho_m}\partial^{s_n}_{\rho_n}\partial^{s_p}_{\rho_p}\sum_{m,n,p}c(m,n,p;\rho_m,\rho_n,\rho_p)z_1^{m+\rho_m}z_2^{n+\rho_n}z_3^{p+\rho_p}|_{\rho_m=\rho_n=\rho_p=0}, \tag{28}$$

with $s_m + s_n + s_p \leq 3$, and $z_1 \equiv \frac{\mu_1\mu_2}{\mu_3^2}$, $z_2 \equiv \frac{\mu_3\mu_4}{\mu_7^2}$, $z_3 \equiv \frac{\mu_7\mu_5^2\mu_6^3}{\mu_0^6}$, needs to be understood. The appearance of $\partial^{s_m}_{\rho_m}\partial^{s_n}_{\rho_n}\partial^{s_p}_{\rho_p}\sum_{m,n,p}$ in (28) is what was referred to earlier on as parametric differentiation of infinite series, something which, as we have explicitly shown above, is not needed in the approach followed in this work.

The Picard-Fuchs equation can be written in the form[10]:

$$\Big(\Delta_z^8 + \sum_{i=1}^{7}\mathbf{B}_i(z)\Delta_z^i\Big)\tilde{\Pi}(z) = 0. \tag{29}$$

The Picard-Fuchs equation in the form written in (29) can alternatively be

---

[4] In equations (20) - (27), use has been made of the following identities:

$$\begin{gathered}\prod_{l=0}^{n-1}\Gamma(x+\tfrac{l}{n}) = (2\pi)^{\frac{n-1}{2}}n^{\frac{1}{2}-nx}\Gamma(nx),\\ \sum_{l=0}^{n-1}\psi(x+\tfrac{l}{n}) - n\psi(nx) = -nlnn,\\ \psi(1) = -\gamma,\ \psi^{(n)} = (-)^{n-1}n!\zeta(n+1).\end{gathered}$$

Using these, one arrives at:

$$\begin{gathered}\psi(\tfrac{1}{3})+\psi(\tfrac{2}{3}) = -2\gamma - ln3^3,\\ \psi(\tfrac{1}{3})-\psi(\tfrac{2}{3}) = -\tfrac{\pi}{\sqrt{3}},\\ \sum_{j=1}^{5}\psi(\tfrac{j}{6}) = -5\gamma - ln6^6\\ \psi'(\tfrac{1}{3})+\psi'(\tfrac{2}{3}) = 8\zeta(2) = \tfrac{4}{3}\pi^2,\\ \psi''(\tfrac{1}{3})+\psi''(\tfrac{2}{3}) = -52\zeta(3),\\ \psi'''(\tfrac{1}{3})+\psi'''(\tfrac{2}{3}) = 480\zeta(4) = \tfrac{16\pi^4}{3},\\ \sum_{j=1}^{5}\psi'(\tfrac{j}{6}) = 35\zeta(2) = \tfrac{35}{6}\pi^2,\\ \sum_{j=1}^{5}\psi''(\tfrac{j}{6}) = -430\zeta(3),\\ \sum_{j=1}^{5}\psi'''(\tfrac{j}{6}) = 7770\zeta(4) = \tfrac{259\pi^4}{3}.\end{gathered}$$

expressed as the following system of eight linear differential equations:

$$\Delta_z \begin{pmatrix} \tilde{\Pi}(z) \\ \Delta_z\tilde{\Pi}(z) \\ (\Delta_z)^2\tilde{\Pi}(z) \\ ... \\ (\Delta_z)^7\tilde{\Pi}(z) \end{pmatrix} =$$

$$\begin{pmatrix} 0 & 1 & 0 & ...0 & 0 \\ 0 & 0 & 1 & ...0 & 0 \\ . & . & . & .... & . \\ 0 & 0 & 0 & ...0 & 1 \\ 0 & -\mathbf{B}_1(z) & -\mathbf{B}_2(z) & ... - \mathbf{B}_6(z) & -\mathbf{B}_7(z) \end{pmatrix} \begin{pmatrix} \tilde{\Pi}(z) \\ \Delta_z\tilde{\Pi}(z) \\ (\Delta_z)^2\tilde{\Pi}(z) \\ ... \\ (\Delta_z)^7\tilde{\Pi}(z) \end{pmatrix} \quad (30)$$

The matrix on the RHS of (30) is usually denoted by $A(z)$.

If the eight solutions, $\{\tilde{\Pi}_{I=1,...,8}\}$, are collected as a column vector $\tilde{\Pi}(z)$, then the *constant*[5] monodromy matrix T for $|z| << 1$ is defined by:

$$\tilde{\Pi}(e^{2\pi i}z) = T\tilde{\Pi}(z). \quad (31)$$

The basis for the space of solutions can be collected as the columns of the "fundamental matrix" $\Phi(z)$ given by:

$$\Phi(z) = S_8(z)z^{R_8}, \quad (32)$$

where $S_8(z)$ and R_8 are 8×8 matrices that single and multiple-valued respectively. Note that $\mathbf{B}_i(0) \neq 0$, which influences the monodromy properties. Also,

$$\Phi(z)_{ij} = \begin{pmatrix} \tilde{\Pi}_1(z) & ... & \tilde{\Pi}_8(z) \\ \Delta_z\tilde{\Pi}_1(z) & ... & \Delta\tilde{\Pi}_8(z) \\ \Delta_z^2\tilde{\Pi}_2(z) & ... & \Delta^2\tilde{\Pi}_8(z) \\ ... & ... & ... \\ \Delta_z^7\tilde{\Pi}_1(z) & ... & \Delta_z^7\tilde{\Pi}_8(z) \end{pmatrix}_{ij} , \quad (33)$$

implying that

$$T = e^{2\pi i R^t}. \quad (34)$$

Now, writing $z^R = e^{Rlnz} = 1 + Rlnz + R^2(lnz)^2 + ...$, and further noting that there are no terms of order higher than $(lnz)^4$ in $\tilde{\Pi}(z)$ obtained above, implies that the matrix R must satisfy the property: $R^m = 0,\ m = 5, ...\infty$. Hence, $T = e^{2\pi i R^t} = 1 + 2\pi i R^t + \frac{(2\pi i)^2}{2}(R^t)^2 + \frac{(2\pi i)^3}{6}(R^t)^3 + \frac{(2\pi i)^4}{24}(R^t)^4$. Irrespective of whether or not the distinct eigenvalues of $A(0)$ differ by integers, one has to evaluate $e^{2\pi i A(0)}$. The eigenvalues of $A(0)$ of (A1), are $0^5, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}$, and hence five of the eight eigenvalues differ by an integer (0).

---

[5] This thus implies that both $\tilde{\Pi}$ and Π, have the *same* monodromy matrix.

Now, the Picard-Fuchs equation (13) can be rewritten in the form (29), with the following values of B_i's:

$$\begin{aligned}
\mathbf{B}_{1,2} &= 0 \\
\mathbf{B}_3 &= \frac{5z}{324(1+z)} \\
\mathbf{B}_4 &= \frac{137}{648(1+z)} \\
\mathbf{B}_5 &= \frac{(\frac{25}{24}z - \frac{1}{9})}{(1+z)} \\
\mathbf{B}_6 &= \frac{(26+85z)}{36(1+z)} \\
\mathbf{B}_7 &= -\frac{(3-5z)}{2(1+z)}.
\end{aligned} \tag{35}$$

Under the change of basis $\tilde{\Pi}(z) \to \tilde{\Pi}'(z) = M^{-1}\tilde{\Pi}(z)$, and writing $\tilde{\Pi}_j(z) = \sum_{i=0}^4 (lnz)^i q_{ij}(z)$ (See [8] and the appendix), one sees that

$$\begin{aligned}
&\tilde{\Pi}'_j(z) = \sum_{i=0}^{4}(lnz)^i q'_{ij}(z), \\
&q'(z) = q(z)(M^{-1})^t, \\
&\Phi'(z)\Phi(z)(M^{-1})^t,\ S'(z) = S(z)(M^{-1})^t,\ R' = M^t R (M^{-1})^t.
\end{aligned} \tag{36}$$

By choosing M such that $S'(0) = \mathbf{1}_{24}$, one gets

$$T(0) = M(e^{2i\pi A(0)})^t M^{-1}. \tag{37}$$

We evaluate $(M^{-1})^t$ and $e^{2\pi i A(0)}$ in the appendix. From the form of $(M^{-1})^t$, one gets:

$$\begin{aligned}
M = \{\{&1, 0, 0, \frac{-6\ (-16 + 16\, q_{00} - 29499\, q_{10})}{1617589}, \frac{5586 - 5586\, q_{00} + 795503\, q_{10}}{14558301}, \\
&\frac{-4\ (-110 + 110\, q_{00} - 607\, q_{10})}{4852767}, \frac{-5\ (-2052 + 2052\, q_{00} + 665123\, q_{10})}{131024709}, \\
&\frac{16212 - 16212\, q_{00} - 8527877\, q_{10}}{262049418}\}, \{0, 1, 0, \frac{-2\ (88497 + 48\, q_{01} - 88497\, q_{11} + 1030121\, q_{21})}{1617589},
\end{aligned}$$

$$\frac{-795503-5586\,q_{01}+795503\,q_{11}-6851300\,q_{21}}{14558301},\frac{-4\ (607+110\,q_{01}-607\,q_{11}+1710\,q_{21})}{4852767},$$

$$\frac{3325615-10260\,q_{01}-3325615\,q_{11}+25280767\,q_{21}}{131024709},$$

$$\frac{8527877-16212\,q_{01}-8527877\,q_{11}+62539841\,q_{21}}{262049418}\},$$

$$\{0,0,1,\frac{-96\,q_{02}+176994\,q_{12}-1030121\ (-1+2\,q_{22})}{1617589},$$

$$\frac{-5586\,q_{02}+795503\,q_{12}-3425650\ (-1+2\,q_{22})}{14558301},\frac{-4\ (110\,q_{02}-607\,q_{12}+855\ (-1+2\,q_{22}))}{4852767},$$

$$\frac{-20520\,q_{02}-6651230\,q_{12}+25280767\ (-1+2\,q_{22})}{262049418},$$

$$\frac{-32424\,q_{02}-17055754\,q_{12}+62539841\ (-1+2\,q_{22})}{524098836}\},$$

$$\{0,0,0,\frac{-2\ (48\,q_{03}-88497\,q_{13}+1030121\,q_{23})}{1617589},$$

$$\frac{-5586\,q_{03}+795503\,q_{13}-6851300\,q_{23}}{14558301},$$

$$\frac{-4\ (110\,q_{03}-607\,q_{13}+1710\,q_{23})}{4852767},\frac{-10260\,q_{03}-3325615\,q_{13}+25280767\,q_{23}}{131024709},$$

$$\frac{-16212\,q_{03}-8527877\,q_{13}+62539841\,q_{23}}{262049418}\},$$

$$\{0,0,0,\frac{-2\ (48\,q_{04}-88497\,q_{14}+1030121\,q_{24}-4852767\,q_{34})}{1617589},$$

$$\frac{-5586\,q_{04}+795503\,q_{14}-6851300\,q_{24}+349399224\,q_{44}}{14558301},$$

$$\frac{-4\ (110\,q_{04}-607\,q_{14}+1710\,q_{24})}{4852767},\frac{-10260\,q_{04}-3325615\,q_{14}+25280767\,q_{24}}{131024709},$$

$$\frac{-16212\,q_{04}-8527877\,q_{14}+62539841\,q_{24}}{262049418}\},$$

$$\{0,0,0,\frac{-2\ (48\,q_{05}-88497\,q_{15}+1030121\,q_{25}-4852767\,q_{35})}{1617589},$$

$$\frac{-5586\,q_{05}+795503\,q_{15}-6851300\,q_{25}}{14558301},$$

$$\frac{-4\ (110\,q_{05}-607\,q_{15}+1710\,q_{25})}{4852767},\frac{-10260\,q_{05}-3325615\,q_{15}+25280767\,q_{25}}{131024709},$$

$$\frac{-16212\,q_{05}-8527877\,q_{15}+62539841\,q_{25}}{262049418}\},$$

$$\{0,0,0,\frac{-2\ (48\,q_{06}-88497\,q_{16}+1030121\,q_{26}-4852767\,q_{36})}{1617589},$$

$$
\begin{aligned}
&\frac{-5586\,q_{06}+795503\,q_{16}-6851300\,q_{26}}{14558301},\\
&\frac{-4\,(110\,q_{06}-607\,q_{16}+1710\,q_{26})}{4852767},\frac{-10260\,q_{06}-3325615\,q_{16}+25280767\,q_{26}}{131024709},\\
&\frac{-16212\,q_{06}-8527877\,q_{16}+62539841\,q_{26}}{262049418}\},\\
&\{0,0,0,\frac{-2\,(48\,q_{07}-88497\,q_{17}+1030121\,q_{27}-4852767\,q_{37})}{1617589},\\
&\frac{-5586\,q_{07}+795503\,q_{17}-6851300\,q_{27}}{14558301},\\
&\frac{-4\,(110\,q_{07}-607\,q_{17}+1710\,q_{27})}{4852767},\frac{-10260\,q_{07}-3325615\,q_{17}+25280767\,q_{27}}{131024709},\\
&\frac{-16212\,q_{07}-8527877\,q_{17}+62539841\,q_{27}}{262049418}\}\}
\end{aligned}
\tag{38}
$$

The elements q_{ij} of a matrix q, are as explained in the appendix. Using MATHEMATICA, one can actually evaluate T, but the expression is extremely long and complicated and will not be given in this paper.

The monodromy around $z = \infty$ can be evaluated as follows(similar to the way given in [8]). For $|z| >> 1$, one can write:

$$
\tilde{\Pi}_a(z) = \sum_{j=1}^{5} A_{aj}(z)u_j(z),\ a = 0, ..., 7,
\tag{39}
$$

where $u_j(z) = e^{-\frac{j}{6}}$. Now, as $z \rightarrow e^{2i\pi}z$, with obvious meanings to the notation:

$$
T_u(\infty) = \begin{pmatrix} e^{-i\frac{\pi}{3}} & 0 & 0 & 0 & 0 \\ 0 & e^{-\frac{2i\pi}{3}} & 0 & 0 & 0 \\ 0 & 0 & e^{-i\pi} & 0 & 0 \\ 0 & 0 & 0 & e^{-\frac{4i\pi}{3}} & 0 \\ 0 & 0 & 0 & 0 & e^{-\frac{5i\pi}{3}} \end{pmatrix}.
\tag{40}
$$

Now, using

$$
\tilde{\Pi}(z \rightarrow e^{2i\pi}z)|_{z\rightarrow\infty} = \mathcal{A}(z \rightarrow e^{2i\pi}z)T_u(\infty)u(z)|_{z\rightarrow\infty} \equiv T(\infty)\mathcal{A}(z)u(z)|_{z\rightarrow\infty}.
\tag{41}
$$

So, equation (41) is the defining equation for the monodromy matrix around $z \rightarrow \infty$. Note, however, that from the point of view of computations, given that the matrix A is not a square matrix, (41) involves solving 40 equations in 64 variables. The matrix $\mathcal{A}$ is given in appendix B. Unfortunately, MATHEMATICA is not able to handle such a computation, this time. However, it is clear that it is in prinicple, a doable computation.

3 Superpotential Calculation

As done in [11], consider F-theory on an elliptically fibered Calabi-Yau 4-fold X_4 with holomorphic map $\pi : X_4 \rightarrow B_3$ and a 6-divisor D_3 as a section such that $\pi(D_3) = C_2 \subset B_3$. Then for vanishing size of the the elliptic fiber, it was argued in [11] that 5-branes wrapped around D_3 in M-theory on the same X_4 obeying the unit-arithmetic genus condition, $\chi(D_3, \mathcal{O}_{D_3}) = 1$, correspond to 3-branes wrapped around C_2 in type IIB, or equivalently F-theory 3-branes wrapped around $C_2 \subset B_3$. It was shown in [11] that only 3-branes contribute to the superpotential in F-theory. As there are no 3-branes in the F-theory dual [3], this implies that no superpotential is generated on the F-theory side. As F-theory 3-branes correspond to Heterotic instantons, one again expects no superpotential to be generated in Heterotic theory on the self-mirror $CY_3(11, 11)$ based on the $\mathcal{N} = 2$ type IIA/Heterotic dual of Ferrara et al where the same self-mirror Calabi-Yau figured on the type IIA side and the self-mirror nature was argued to show that there are no world-sheet or space-time instanton corrections to the classical moduli space.

If the abovementioned triality is correct, then one must be able to show that there is no superpotential generated on type IIA side on the freely-acting antiholomorphic involution of $CY_3(3, 243)$.

On the mirror type IIB side, the W is generated from domain-wall ($\equiv D5$-branes wrapped around supersymmetric 3-cycles $\hookrightarrow CY_3$'s) tention. $W_{IIB} = \int_{C:\partial C=\sum_i D_i} \Omega_3$, D_i's are 2-cycles corresponding to the positions of $D5$-branes or $O5$-planes, i.e., objects carrying $D5$ brane charge. From the world-sheet point of view, the $D5$ branes correspond to disc amplitudes and $O5$-planes correspond to $\mathbf{RP}^2$ amplitudes. As there are no branes in our theory, we need to consider only $\mathbf{RP}^2$ amplitudes. Now, type IIA on a freely acting involution of a Calabi-Yau with no branes or fluxes can still generate a superpotential because it is possible that free involution on type IIA side corresponds to orientifold planes in the mirror type IIB side, which can generate a superpotential. The same can also be studied using localization techniques in unoriented closed string enumerative geometry [12]. Consider an orientation-reversing diffeomorphism $\sigma : \Sigma \rightarrow \Sigma$, an antiholomorhpic involution on the Calabi-Yau X $I : X \rightarrow X$ and an equivariant map $f : \Sigma \rightarrow X$ [satisfying $f \circ \sigma = I \circ f$]), then the quotient spaces in $\tilde{f} : \frac{\Sigma}{<\sigma>} \rightarrow \frac{X}{I}$ possesses a dianalytic structure.

In the unoriented theory, one then has to sum over holomorphic and antiholomorphic instantons. For connected $\frac{\Sigma}{<\sigma>}$, the two contributions are the same; hence sufficient to consider only equivariant holomorphic maps. One constructs one-dimensional torus action, T, on X compatible with I with isolated fixed points. The action T induces an action on the moduli space of equivariant holomorphic maps, and one then evaluates the localized contributions from the fixed points, using an equivariant version of the the Atiyah-Bott formula, much on the lines of Kontsevich's work. For a Calabi-Yau 3-fold, the virtual cycle "$[\bar{M}_{g,0}(X,\beta)]^{virt}$" is zero-dimensional, and one has to evaluate $\int_{\Xi_s^{virt}} \frac{1}{e_T(N^{virt}_{\Xi_s})}$, where $\Xi_s \equiv$ is the fixed locus in the moduli space of symmetric holomorphic maps, and one sees that one gets a match with similar calculations based on large N dualities and mirror symmetry

For $\int d^2\theta W_{LG}$ to be invariant under $\Omega.\omega$, given that the measure is reflected under Ω, $\omega : W_{LG} \to -W_{LG}$.

away from the orbifold singularities: Promoting the action of ω to the one on the chiral superfields:

$$\omega : (\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_4, \mathcal{X}_5, \mathcal{X}_6, \mathcal{X}_0) \to (\bar{\mathcal{X}}_2, -\bar{\mathcal{X}}_1, \bar{\mathcal{X}}_4, \bar{\mathcal{X}}_5, \bar{\mathcal{X}}_6, \bar{\mathcal{X}}_0), \tag{42}$$

and using $Re(Y_i) = |\mathcal{X}_i|^2$, one gets the following action of ω on the twisted chiral superfields Y_i's:

$$\omega : Y_1 \to Y_2 + i\pi,\ Y_2 \to Y_1 + i\pi;\ Y_{0,4,5,6} \to Y_{0,4,5,6} + i\pi. \tag{43}$$

The action of ω on $Y_{4,5,6,0}$ implies that ω acts without fixed points even on the twisted chiral superfields, further implying that there are no orientifold fixed planes, and *hence no superpotential is generated on the type IIA side away from the orbifold singularities.*

after singularity resolution: Writing $W = \prod_{i=0}^{7} a_{e_0,\ldots,7}\mathcal{X}_i^{e_i}$ with the requirements that $\vec{l^{(a)}} \cdot \vec{e} = 0$ for $a = 1, 2, 3$ and $e_i \leq 1$[13][6], one sees that $\mathcal{X}_0 \prod_{i=1}^{7} \mathcal{X}_i$ is an allowed term in the superpotential. A valid antiholomorphic involution this time can be:

$$\omega : (\mathcal{X}_0, \mathcal{X}_1, \mathcal{X}_2, \mathcal{X}_3, \mathcal{X}_4, \mathcal{X}_5, \mathcal{X}_6, \mathcal{X}_7) \to (\bar{\mathcal{X}}_0, \bar{\mathcal{X}}_2, -\bar{\mathcal{X}}_1, -\bar{\mathcal{X}}_3, \bar{\mathcal{X}}_4, \bar{\mathcal{X}}_5, \bar{\mathcal{X}}_6, \bar{\mathcal{X}}_7). \tag{44}$$

This on the mirror LG side again implies that one will have free actions w.r.t. $Y_{0,3,4,5,6,7}$ implying there can be no orientifold planes and *no superpotential (is likely to be) generated even after singularity resolution.*

4 Action of the antiholomorphic involution on M theory compactified on $CY_3(3,243) \times S^1$

The M-theory uplift of the type IIA side of the $\mathcal{N} = 1$ Heterotic/type IIA dual pair of [5], as obtained in [3], involves the 'barely G_2-manifold'

---

[6] We thank A.Klemm for bringing [13] to our attention.

$\frac{CY_3(3,243)\times S^1}{\mathbf{Z}_2}$. In this section we consider the $D=11$ supergravity limit of M-theory and construct the $\mathcal{N}=1, D=4$ supergravity action, and evaluate the Kähler potential for the same.

The effect of the $\mathbf{Z}_2$ involution that reflects the S^1, $H^{1,1}(CY_3)$ and takes $H^{p,q}(CY_3)$ to $H^{q,p}(CY_3)$ for $p+q=3$, where the CY_3 is the one that figures in $\frac{CY_3\times S^1}{\mathbf{Z}_2}$, at the level of $D=11$ supergravity can be obtained by first compactifying the same on an S^1, then on CY_3 (following [4]) and eventually modding out the action by the abovementioned $\mathbf{Z}_2$ action.

The $D=11$ supergravity action of Cremmer et al is:

$$\mathcal{L}_{11} = -\frac{1}{2}e_{11}R_{11} - \frac{1}{48}(G_{MNPQ})^2 + \frac{\sqrt{2}}{(12)^4}\epsilon^{M_0\ldots M_{10}}G_{M_0\ldots M_3}G_{M_4\ldots M_7}C^{11}_{M_8M_9M_{10}}, \tag{45}$$

which after dimensional reduction on an S^1, gives:

$$\begin{aligned}\mathcal{L}_{10} = &-\frac{1}{2}e_{10}R_{10} - \frac{1}{8}e_{10}\phi^{\frac{9}{4}}F^2_{mn} - \frac{9}{16}e_{10}(\partial_m ln\phi)^2 - \frac{1}{48}e_{10}\phi^{\frac{3}{4}}(F_{mnpq}+6F_{[mn}B_{pq]})^2\\ &-\frac{1}{12}e_{10}\phi^{-\frac{3}{2}}H^2_{mnp} + \frac{\sqrt{2}}{(48)^2}\epsilon^{m_0\ldots m_9}(F_{m_0\ldots m_3}+6F_{m_0m_1}B_{m_2m_3})F_{m_4\ldots m_7}B_{m_8m_9},\end{aligned} \tag{46}$$

where

$$\begin{aligned}&G_{MNPQ} = \partial_{[M}C_{NPQ]}\\ &F_{mnpq} = 4\partial_{[m}C_{npq]};\ B_{mn} = C_{mn10};\ H_{mnp} = 3\partial_{[m}B_{np]};\ F_{mn} = 2\partial_{[m}A_{n]};\\ &C_{mnp} = A_{mnp} + 3A_{[m}B_{np]},\end{aligned} \tag{47}$$

and

$$\begin{aligned}&e_{11}\,{}^{A}_{M} = \begin{pmatrix} e_{10}\,{}^{a}_{m} & \phi A_M \\ 0 & \phi \end{pmatrix}\\ &A, M = 0, \ldots, 10;\ a, m = 0, \ldots, 9.\end{aligned} \tag{48}$$

After compactifying on a CY_3, one gets the following Lagrangian density:

$$\mathcal{L}_4 = \mathcal{L}_4^{\text{grav}+H^0} + \mathcal{L}_4^{H^2} + \mathcal{L}_4^{H^3}, \tag{49}$$

where after a suitable Weyl scaling:

$$\mathcal{L}_4^{\text{grav}} = e\Big[-\frac{R}{2} - \frac{9}{16}\Big(\frac{\partial_\mu\phi}{\phi}\Big)^2 + G_{\alpha\bar{\beta}}\partial_\mu Z_\alpha\partial^\mu\bar{Z}^{\bar{\beta}} - \partial_\mu M^A\partial^\mu M^B\Big(\frac{1}{2}G_{AB} + \frac{9}{4}\frac{\int e^A\wedge J\wedge J\int e^B\wedge J\wedge J}{(\int J\wedge J\wedge J)^2}\Big)\Big], \tag{50}$$

where $A, B = 1, ..., h^{1,1}(CY_3)$, $\alpha, \beta = 1, ..., h^{2,1}(CY_3)$,

$$\begin{aligned} G_{AB} &= \frac{i \int e^A \wedge * e^B}{2\mathcal{V}}, \\ \mathcal{V} &= \frac{1}{3!} \int J \wedge J \wedge J, \\ G_{\alpha\bar{\beta}} &= -\frac{i \int b_\alpha \wedge \bar{b}_\beta}{\mathcal{V}}, \end{aligned} \tag{51}$$

where

$$\bar{b}_{\alpha\ ij} = \frac{i}{||\Omega||^2} \Omega_i^{\overline{lk}} \bar{\Phi}_{\alpha\ \overline{lk}j}, \tag{52}$$

Φ being a (2,1) form, and the $h^{1,1}$ moduli M^A entering in the variation of the metric with mixed indices and the $h^{2,1}$ moduli Z_α entering in the variation of the metric with same indices. Writing (50) in terms of v^A defined via: $M^A = \sqrt{2} v^A \phi^{-\frac{3}{4}}$, one gets:

$$\mathcal{L}_4^{\rm grav} = e\Big[-\frac{R}{2} - \frac{G_{AB}}{2} \partial_\mu v^A \partial^\mu v^B - \frac{1}{4} \frac{\partial_\mu (\mathcal{V}\phi^{-3})^2}{(\mathcal{V}\phi^{-3})^2} + G_{\alpha\bar{\beta}} \partial_\mu Z^\alpha \partial^\mu \bar{Z}^\beta\Big],$$

Under the freely-acting antiholomorphic involution, the $h^{1,1}$-moduli M^A/v^A get projected out, G_{AB} is even, and A_μ^A gets projected out. Thus, one gets:

$$\mathcal{L}_4^{\rm grav}/\mathbf{Z}_2 = e\Big[-\frac{R}{2} - \frac{1}{4} \frac{(\partial_\mu (\mathcal{V}\phi^{-3}))^2}{(\mathcal{V}\phi^{-3})^2} + G_{\alpha\bar{\beta}} \partial_\mu Z^\alpha \partial^\mu \bar{Z}^\beta\Big]. \tag{53}$$

Additionally, after a Weyl scaling:

$$\mathcal{L}_4^{H^0(A_\mu)} = -\frac{\sqrt{2}^3}{8} e\mathcal{V}(v) F_{\mu\nu}^2 \tag{54}$$

$$\mathcal{L}_4^{H^0(A_{\mu\nu\rho})} = -\frac{1}{48} e\mathcal{V}^3(v) \phi^{\frac{3}{4}} (F_{\mu\nu\rho} + 6F_{[\mu\nu} B_{\rho\sigma]})^2, \tag{55}$$

Thus, under the $\mathbf{Z}_2$-involution, $\mathcal{L}_4^{H^0(A_\mu)}$ and $\mathcal{L}_4^{H^0(A_{\mu\nu\rho})}$ get projected out. The term $\mathcal{L}_4^{H^0(H_{\mu\nu\rho},\phi)}$ will be considered in conjunction with $\mathcal{L}_4^{H^3}$ later.

Now, the H^2-sector Lagragian density is given by:

$$\begin{aligned} \mathcal{L}_4^{H^2} &= e\Big[-\frac{1}{2} G_{AB} \partial_\mu W^A \partial^\mu W^B + \frac{1}{\sqrt{2}} F_{\mu\nu}^2 \Big(-\frac{1}{12} \int J \wedge J \wedge J \\ &+ \frac{1}{2} \Big[\int e^A \wedge e^B \wedge J - \frac{3}{2} \frac{\int e^A \wedge J \wedge J \int e^B \wedge J \wedge J}{\int J \wedge J \wedge J} \Big) a^A a^B\Big] \end{aligned}$$

$$
\begin{aligned}
&+\frac{1}{\sqrt{2}}\Big(\int e^A\wedge e^B\wedge J-\frac{3}{2}\frac{\int e^A\wedge J\wedge J\int e^B\wedge J\wedge J}{\int J\wedge J\wedge J}f^A_{\mu\nu}a^Bf^{\mu\nu}\\
&+\frac{\sqrt{2}}{4}\Big(\int e^A\wedge e^B\wedge J-\frac{3}{2}\frac{\int e^A\wedge J\wedge J\int e^B\wedge J\wedge J}{\int J\wedge J\wedge J}\Big)f^A_{\mu\nu}f^{\mu\nu\ B}\Big]\\
&+i\Big[\frac{\sqrt{2}}{12}a^Aa^Ba^C\tilde{F}_{\mu\nu}F^{\mu\nu}\int e^A\wedge e^B\wedge e^C+\frac{\sqrt{2}}{8}a^Aa^B\tilde{F}_{\mu\nu}f^{C\ \mu\nu}\int e^A\wedge e^B\wedge e^C\\
&+\frac{\sqrt{2}}{8}a^Aa^BF_{\mu\nu}\tilde{F}^C_{\mu\nu}\int e^A\wedge e^B\wedge e^C+\frac{\sqrt{2}}{4}a^Af^B_{\mu\nu}\tilde{f}^{C\ \mu\nu}\int e^A\wedge e^B\wedge e^C\Big], \qquad (56)
\end{aligned}
$$

where $W^A\equiv a^A+iv^A$ and, e.g., $e\tilde{F}^{\mu\nu}\equiv\frac{1}{2}\epsilon^{\mu\nu\rho\lambda}F_{\rho\lambda}$. Thus, under the $\mathbf{Z}_2$-involution, only $-\frac{1}{2}eG_{AB}\partial_\mu a^A\partial^\mu a^B$ in $\mathcal{L}_4^{H^2}$, survives, i.e.,

$$
\mathcal{L}_4^{H^2}/\mathbf{Z}_2=-\frac{1}{2}eG_{AB}\partial_\mu a^A\partial^\mu a^B. \qquad (57)
$$

Now, using the notations of [4],

$$
\begin{aligned}
&\mathcal{L}_4^{H^3}+\mathcal{L}_4^{H^0}=-e\frac{1}{4\tilde{\phi}^2}|\partial_\mu\tilde{\phi}+i\partial_\mu D-\frac{1}{4}\partial_\mu[(\Psi+\bar{\Psi})R^{-1}(\Psi+\bar{\Psi})]+(\Psi+\bar{\Psi})R^{-1}\partial_\mu\Psi\\
&-\frac{1}{4}(\psi+\bar{\psi})R^{-1}\partial_\mu\mathcal{N}R^{-1}(\Psi+\bar{\Psi})|^2\\
&=-\frac{|\partial_\mu S+(\Psi+\bar{\Psi})R^{-1}\partial_\mu\Psi-\frac{1}{4}(\Psi+\bar{\Psi})R^{-1}\partial_\mu\mathcal{N}R^{-1}(\Psi+\bar{\Psi})|^2}{[S+\bar{S}+\frac{1}{2}(\Psi+\bar{\Psi})R^{-1}(\Psi+\bar{\Psi})]^2}\\
&+\frac{[\partial_\mu\Psi-\frac{1}{2}(\psi+\bar{\Psi})R^{-1}\partial_\mu\mathcal{N}]R^{-1}[\partial_\mu\bar{\Psi}-\frac{1}{2}\partial_\mu\bar{\mathcal{N}}R^{-1}(\Psi+\bar{\Psi})]}{[(S+\bar{S})+\frac{1}{2}(\Psi+\bar{\Psi})R^{-1}(\Psi+\bar{\Psi})]}, \qquad (58)
\end{aligned}
$$

where $S\equiv\tilde{\phi}+iD-\frac{1}{4}(\Psi+\bar{\Psi})R^{-1}(\psi+\bar{\Psi})$, D being a Lagrange multiplier, $\tilde{\phi}\equiv\sqrt{2}\mathcal{V}(v)\phi^{-3}$, $\Psi_{I(\equiv0,1,\ldots,h^{2,1})}$ appearing in the expansion of the real 3-form A_{mnp} in a canonical basis of H^3, and

$$
\begin{aligned}
&R_{IJ}\equiv Re[\mathcal{N}_{IJ}],\ \mathcal{N}_{IJ}\equiv\frac{1}{4}\bar{F}_{IJ}-\frac{(NZ)_I(NZ)_J}{(ZNZ)},\\
&(R^{-1})^{IJ}=2\Big(N^{-1}(\mathbf{1}-\bar{K}\bar{Z}-KZ)\Big)^{IJ}, \qquad (59)
\end{aligned}
$$

where Z^I and iF_I are the period integrals, $N_{IJ}=\frac{1}{4}(F_{IJ}+\bar{F}_{IJ})$, and $K_I\equiv\frac{\int\Omega_I\wedge\bar{\Omega}}{\int\Omega\wedge\bar{\Omega}}$, with $\Omega_I\equiv\frac{\partial\Omega}{\partial Z^I}$. Here, it is assumed that the holomorphic 3-form Ω is expaneded in a canonical cohomology basis (α_I,β^I) satisfying

$$
\int_{A^J}\alpha_I=\int\alpha_I\wedge\beta^J=-\int_{B_I}\beta^J=-\int\beta^J\wedge\alpha_I=\delta_I^J, \qquad (60)
$$

with (A^I, B_I) being the dual homology basis. The period integrals are then defined to be: $Z^I = \int_{A^I} \Omega$ and $iF_I = \int_{B_I} \Omega$. Hence,

$$\Omega = Z^I \alpha_I + iF_I \beta^I. \tag{61}$$

For the $\mathcal{N} = 1$ case, we work in the large volume limit of the Calabi-Yau. In this limit, one gets:

$$\mathcal{L}^{\text{grav}+H^0+H^2+H^3}/\mathbf{Z}_2 = e\Big[-\frac{R}{2} - G_{AB}\partial_\mu a^A \partial^\mu a^B + G_{\alpha\bar{\beta}}\partial_\mu Z^\alpha \partial^\mu Z^{\bar{\beta}} - \frac{1}{2}\frac{(\partial_\mu\tilde{\phi})^2}{\tilde{\phi}^2}\Big]. \tag{62}$$

Hence, one gets for the $\mathcal{N} = 1$ Kähler potential $K_{\mathcal{N}=1}$:

$$K_{\mathcal{N}=1} = K[a^A, Z^\alpha] + \frac{1}{2} ln[\tilde{\phi}]. \tag{63}$$

At the $\mathcal{N} = 2$ level, if there were decoupling between the fields of the H^3-sector from the other fields of the other sectors, the Kähler potential would be consisting of $ln\Big[S + \bar{S} + \frac{1}{2}(\Psi + \bar{\Psi})R^{-1}(\Psi + \bar{\psi})\Big]$,[7] from the H^3-sector. From the definitin of S above, one sees that:

$$S + \bar{S} + \frac{1}{2}(\Psi + \bar{\Psi})R^{-1}(\Psi + \bar{\Psi}) = 2\tilde{\phi}. \tag{64}$$

This partially explains the appearance of $ln[\tilde{\phi}]$ in $K_{\mathcal{N}=1}$.

5 Action of Antiholomorphic Involution on the Periods - a Conjecture

Given the action of an antiholomorphic involution on the cohomology, it is in general quite non-trivial to figure out the action of the involution on the period integrals using the canonical (co)homology basis of (60). We now discuss a reasonable guess for the same. From (64), one sees that the RHS is reflected under the antiholomorphic involution discussed towards the beginning of this section. We now conjecture that on the LHS, this would imply that

$$\begin{aligned} &S \to -S,\ (\Psi + \bar{\Psi})^2 \to (\Psi + \bar{\Psi})^2, \\ &R_{IJ} \to -R_{IJ}. \end{aligned} \tag{65}$$

---

[7] This however assumes that $\frac{\partial \bar{\mathcal{N}}_{IJ}}{\partial z^K} = 0 \leftrightarrow \frac{1}{4}F_{IJK} - \frac{(\frac{1}{4}F_{IRK}\bar{z}^R(N\bar{Z})_J + \frac{1}{4}(N\bar{Z}_I F_{JLK}\bar{Z}^L)}{(\bar{Z}N\bar{Z})} + \frac{(N\bar{Z})_I(N\bar{Z})_J}{(2\bar{Z}N\bar{Z})^2}\bar{Z}^P\bar{Z}^Q F_{PQK} = 0$

We further conjecture that $R_{IJ} \to -R_{IJ}$ is realized by

$$\begin{aligned} & Z^I \to -\bar{Z}^I,\ F_I \to \bar{F}_I \\ & \alpha_I \to -\alpha_I,\ \beta^I \to \beta^I. \end{aligned} \tag{66}$$

One should note that given that the antiholomorphic involution is orientation reversing, the intersection form $\int \alpha_I \wedge \beta^J$ is also reflected.[8] This fact, e.g., can be explicitly seen in the real basis of $H^3(T^6, \mathbf{Z})$[14]:

$$\begin{aligned} & \alpha_0 = dx^1 \wedge dx^2 \wedge dx^3;\ \alpha_{ij} = \frac{1}{2}\epsilon_{ilm} dx^l \wedge dx^m \wedge dy^j (1 \leq i, j \leq 3), \\ & \beta_0 = dy^1 \wedge dy^2 \wedge dy^3;\ \beta^{ij} = -\frac{1}{2}\epsilon_{jlm} dy^l \wedge dy^m \wedge dx^i (1 \leq i, j \leq 3). \end{aligned} \tag{67}$$

where $x^i, y^i, i = 1, 2, 3$ are the six real coordinates on the T^6, using which one can construct the holomorphic one-form: $dz^i = dx^i + \tau^{ij} dy^j$, τ^{ij} being the period matrix for the torus. Hence, under the antiholomorphic involution: $(x^i, y^i, \tau^{ij}) \to (x^i, -y^i, -\bar{\tau}^{ij})$, the intersection matrix changes sign because α_{ij} and β_0 change signs.[9] The holomorphic 3-form Ω is then given by:

$$\Omega = dz^1 \wedge dz^2 \wedge dz^3 = \alpha_0 - \beta^{ij}(cof\tau)_{ij} + \alpha_{ij}\tau^{ij} + \beta_0(det\tau), \tag{68}$$

which on comparison with:

$$\Omega = Z^I \alpha_I + i F_I \beta^I, \tag{69}$$

for inhomogenous coordinates ($I = 0, 1, ..., h^{2,1}$ and $h^{2,1}(T^6) = 9$), implies that

$$\begin{aligned} & (\alpha_0, \beta^{ij}) \sim \beta^I,\ (\beta_0, \alpha_{ij}) \sim \alpha_I; \\ & (1, -(cof\tau)_{ij}) \sim F_I,\ (det\tau, \tau^{ij}) \sim Z^I. \end{aligned} \tag{70}$$

One thus sees that the conjecture (66) is satisfied. Further, for a real canonical cohomology basis (α_I, β^I), $\bar{\Omega}$ will be given by $\bar{\Omega} = \bar{Z}^I \alpha_I - i\bar{F}_I \beta^I$. One sees that (66) is consistent with complex conjugation of the holomorphic 3-form Ω.

Further, as a simple toy example, one can consider the mirror to the quintic, for which $h^{2,1} = 1$. If (p, q) forms are denoted by $X^{p,q}$, then one choice of real canonical cohomology basis satisfying (60) is

$$\begin{aligned} & a_0 = \frac{i}{\sqrt{2}}(\Omega - \bar{\Omega}),\ a_1 = \frac{i}{\sqrt{2}}(X^{2,1} - X^{1,2}); \\ & b^0 = \frac{1}{\sqrt{2}}(\Omega + \bar{\Omega}),\ b^1 = \frac{1}{\sqrt{2}}(X^{2,1} + X^{1,2}). \end{aligned} \tag{71}$$

---

[8] We are grateful to D.Joyce for pointing this out to us.

[9] We thank M.Schulz for pointing this out to us.

One sees that that indeed, (a_0, a_1) under the aformentioned antiholomorphic involution goes to $-(a_0, a_1)$ and (b^0, b^1) goes to (b^0, b^1)

6 Conclusion and Discussions

We obtained the Meijer basis of solution to the Picard-Fuchs equation for the Landau-Ginsburg model corresponding to $CY_3(3,243)$ *after* the resolution of the orbifold singularities of the degree-24 Fermat hypersurface in $\mathbf{WCP}^4[1,1,2,8,12]$, in the large *and* small complex structure limits, getting the *ln*-terms without resorting to parametric differentiations of infinite series. We also discussed in detail the evaluation of the monodromy matrix in the large complex structure limit. We also considered the action of an antiholomorphic involution on $D = 11$ supergravity compactified on $CY_3(3,243) \times S^1$, and evaluated the form of the $\mathcal{N} = 1$ Kähler potential. In the process, we also gave a conjecture on the action of the involution on the periods of $CY_3(3,243)$, given its action on the cohomology of the same. We verified the conjecture for T^6 for the periods and cohomology basis, and for the mirror to the quintic for the cohomology basis. Finally, we showed that no superpotential is generated in type IIA and hence M-theory sides using mirror symmetry, *after* the resolution of the orbifold singularities associated with the Fermat hypersurface whose blow up gives $CY_3(3,243)$.

For future work, it will be nice to explicitly work out the Kähler potential, or equivalently the periods for the "(12,12)" $E_8 \times E_8$ Heterotic model, analogous to the "(14,10)" model of [15, 16] and perhaps using the conjecture that both models lie in the same moduli space[10] [17], so as to be able to compare with the Kähler potential of type IIA on $CY_3(3,243)$ or equivalently $D = 11$ supergravity on $\frac{CY_3(3,243)\times S^1}{\mathbf{Z}_2}$. Further, it will be interesting to see whether or not the conjectured $\mathcal{N} = 1$ triality holds up when one turns on fluxes[18] - issues of the right choice of non-Kähler manifolds, flux-induced superpotentials, etc will become relevant for investigation.

Acknowledgements

We are extremely grateful to A.Klemm and D.Joyce for very useful discussions and clarifications, T.Mohaupt and M.Schulz for useful communications,

---

[10] We are grateful to T.Mohaupt for pointing this out to us and for bringing [15] to our attention

and to D.Lüst for going through a preliminary version of the manuscript. The research work is supported by the Alexander von Humboldt foundation.

A Monodromy around $z = 0$

In this appendix, we give the details relevant to the evaluation of the matrix M that is essential for the evaluation of the monodromy matrix T. MATHEMATICA has been heavily used in this appendix. We first give the exponentiation of the relevant matrix $A(0)$ followed by the value of $z^{A(0)}$, again necessary for evaluation of M and hence T.

The matrix $A(0)$ is given by:

$$A(0) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & \frac{1}{9} & -\frac{13}{18} & \frac{3}{2} \end{pmatrix} \tag{A1}$$

Using MATHEMATICA, one then can evaluate the "matrix exponent" involving $A(0)$. The expression given below, has been written in the MATHEMATICA notebook format (by cutting and pasting the "TerXForm" of the output).

$e^{2\pi i A(0)} = \{\{1, 2\,i\,\pi, -2\,\pi^2, \frac{-4\,i}{3}\,\pi^3, \frac{2\,\pi^4}{3}, -\left(\frac{31001}{32}\right) + \frac{729}{32\,e^{\frac{2\,i}{3}\,\pi}} + 1458\,e^{\frac{2\,i}{3}\,\pi} -$
$\frac{5971\,i}{8}\,\pi + \frac{865\,\pi^2}{4} + \frac{115\,i}{3}\,\pi^3 - \frac{13\,\pi^4}{3}, \frac{93213}{32} - \frac{3645}{32\,e^{\frac{2\,i}{3}\,\pi}} - 5103\,e^{\frac{2\,i}{3}\,\pi} + \frac{19215\,i}{8}\,\pi - \frac{2637\,\pi^2}{4} -$
$105\,i\,\pi^3 + 9\,\pi^4, -\left(\frac{35307}{16}\right) + \frac{2187}{16\,e^{\frac{2\,i}{3}\,\pi}} + 4374\,e^{\frac{2\,i}{3}\,\pi} - \frac{7785\,i}{4}\,\pi + \frac{1035\,\pi^2}{2} + 78\,i\,\pi^3 - 6\,\pi^4\},$
$\{0, 1, 2\,i\,\pi, -2\,\pi^2, \frac{-4\,i}{3}\,\pi^3, 128 + \frac{243}{16\,e^{\frac{2\,i}{3}\,\pi}} + 486\,e^{\frac{2\,i}{3}\,\pi} - \frac{i}{18}\,\left(\frac{-53739\,i}{8} + \frac{7785\,\pi}{2} + 1035\,i\,\pi^2 - 156\,\pi^3\right),$
$-576 - \frac{1215}{16\,e^{\frac{2\,i}{3}\,\pi}} - 1701\,e^{\frac{2\,i}{3}\,\pi} - \frac{i}{18}\,\left(\frac{172935\,i}{8} - \frac{23733\,\pi}{2} - 2835\,i\,\pi^2 + 324\,\pi^3\right), 576 +$
$\frac{729}{8\,e^{\frac{2\,i}{3}\,\pi}} + 1458\,e^{\frac{2\,i}{3}\,\pi} - \frac{i}{18}\,\left(\frac{-70065\,i}{4} + 9315\,\pi + 2106\,i\,\pi^2 - 216\,\pi^3\right)\},$
$\{0, 0, 1, 2\,i\,\pi, -2\,\pi^2, 64 + \frac{81}{8\,e^{\frac{2\,i}{3}\,\pi}} + 162\,e^{\frac{2\,i}{3}\,\pi} + \frac{-\left(\frac{70065}{2}\right) - 18630\,i\,\pi + 4212\,\pi^2}{324}, -288 -$
$\frac{405}{8\,e^{\frac{2\,i}{3}\,\pi}} - 567\,e^{\frac{2\,i}{3}\,\pi} + \frac{\frac{213597}{2} + 51030\,i\,\pi - 8748\,\pi^2}{324}, 288 + \frac{243}{4\,e^{\frac{2\,i}{3}\,\pi}} + 486\,e^{\frac{2\,i}{3}\,\pi} + \frac{-83835 - 37908\,i\,\pi + 5832\,\pi^2}{324}\},$
$\{0, 0, 0, 1, 2\,i\,\pi, 32 + \frac{27}{4\,e^{\frac{2\,i}{3}\,\pi}} + 54\,e^{\frac{2\,i}{3}\,\pi} + \frac{i}{5832}\,(167670\,i - 75816\,\pi), -144 - \frac{135}{4\,e^{\frac{2\,i}{3}\,\pi}} -$
$189\,e^{\frac{2\,i}{3}\,\pi} + \frac{i}{5832}\,(-459270\,i + 157464\,\pi), 144 + \frac{81}{2\,e^{\frac{2\,i}{3}\,\pi}} + 162\,e^{\frac{2\,i}{3}\,\pi} + \frac{i}{5832}\,(341172\,i - 104976\,\pi)\},$

$$\{0,0,0,0,1,\tfrac{19}{2}+\tfrac{9}{2\,e^{\frac{2\,i}{3}\,\pi}}+18\,e^{\frac{2\,i}{3}\,\pi},-\left(\tfrac{117}{2}\right)-\tfrac{45}{2\,e^{\frac{2\,i}{3}\,\pi}}-63\,e^{\frac{2\,i}{3}\,\pi},63+\tfrac{27}{e^{\frac{2\,i}{3}\,\pi}}+54\,e^{\frac{2\,i}{3}\,\pi}\},$$
$$\{0,0,0,0,0,8+\frac{3}{e^{\frac{2\,i}{3}\,\pi}}+6\,e^{\frac{2\,i}{3}\,\pi},-36-\frac{15}{e^{\frac{2\,i}{3}\,\pi}}-21\,e^{\frac{2\,i}{3}\,\pi},36+\frac{18}{e^{\frac{2\,i}{3}\,\pi}}+18\,e^{\frac{2\,i}{3}\,\pi}\},$$
$$\{0,0,0,0,0,4+\frac{2}{e^{\frac{2\,i}{3}\,\pi}}+2\,e^{\frac{2\,i}{3}\,\pi},-18-\frac{10}{e^{\frac{2\,i}{3}\,\pi}}-7\,e^{\frac{2\,i}{3}\,\pi},18+\frac{12}{e^{\frac{2\,i}{3}\,\pi}}+6\,e^{\frac{2\,i}{3}\,\pi}\},$$
$$\{0,0,0,0,0,2+\frac{4}{3\,e^{\frac{2\,i}{3}\,\pi}}+\frac{2\,e^{\frac{2\,i}{3}\,\pi}}{3},-9-\frac{20}{3\,e^{\frac{2\,i}{3}\,\pi}}-\frac{7\,e^{\frac{2\,i}{3}\,\pi}}{3},9+\frac{8}{e^{\frac{2\,i}{3}\,\pi}}+2\,e^{\frac{2\,i}{3}\,\pi}\}\}$$

Writing the solution vector $\tilde{\Pi}_i$ as $\tilde{\Pi}_i = \sum_{j=0}^{4}(lnz)^j q_{ji}$ (following the notation of [8]), one notes:

$$(\Phi')_i = (\tilde{\Pi}')^t_i = \left(S' z^{A(0)}\right)_{0i} = (lnz)^j q'_{ji}. \tag{A2}$$

From (A2), one gets the following:

$$(q'(0))_{ji} = \frac{\delta_{ji}}{j!},\ 0 \le (i,j) \le 4. \tag{A3}$$

For $5 \le i \le 7$, consider. e.g., $i = 5$. Then from the expression for $z^{A(0)}$ above,

$$\begin{aligned}
&\sum_{j=0}^{4}(q')_{j5}(lnz)^j = (S')_{00}[f_{05}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + \sum_{n=1}^{4} c_n^{(05}(lnz)^n] + (S')_{01}[f_{15}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + \sum_{n=1}^{3} c_n^{(25)}(lnz)^n]\\
&(S')_{02}[f_{25}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + \sum_{n=1}^{2} c_n^{(25}(lnz)^n] + (S')_{03}[f_{35}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + c_1^{(35)}(lnz)]\\
&+(S')_{04}f_{45}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + (S')_{05}f_{55}(z^{\frac{1}{3}}) + (S')_{06}f_{65}(z^{\frac{1}{2}}, z^{\frac{1}{3}}) + (S')_{07}f_{75}(z^{\frac{1}{3}})
\end{aligned} \tag{A4}$$

where the f_{ij}'s and c_n^{ij}'s can be determined from the expression for $z^{A(0}$ given below. From (A4), one gets:

$$\begin{aligned}
(q')_{05} &= \sum_{i=0}^{7}(S')_{0i}f_{i5}\\
(q')_{15} &= \sum_{i=0}^{3}(S')_{0i}c_1^{i5}\\
(q')_{25} &= \sum_{i=0}^{2}(S')_{0i}c_2^{i5}\\
(q')_{35} &= \sum_{i=0}^{1}(S')_{0i}c_3^{i5}\\
(q')_{45} &= (S')_{00}c_4^{05}.
\end{aligned} \tag{A5}$$

From (A5), one gets:

$$\begin{aligned} &(q'(0))_{0i} = f_{0i}(0), \\ &(q'(0))_{ij} = c_i^{0j},\ 1 \leq i \leq 4,\ 5 \leq j \leq 7. \end{aligned} \tag{A6}$$

Again using the MATHEMATICA notebook format, the value of $z^{A(0)}$, as evaluated by MATHEMATICA is given by:

$$z^{A(0)} = e^{A(0)ln(z)} = \{\{1, \log(z), \frac{\log(z)^2}{2}, \frac{\log(z)^3}{6}, \frac{\log(z)^4}{24},$$

$$\frac{3\left(-39193+46656\,z^{\frac{1}{3}}-8192\,\sqrt{z}+729\,z^{\frac{2}{3}}\right)-35826\ \log(z)-5190\log(z)^2-460\log(z)^3-26\log(z)^4}{96},$$

$$\frac{-3\left(-43359+54432\,z^{\frac{1}{3}}-12288\,\sqrt{z}+1215\,z^{\frac{2}{3}}-12810\ \log(z)-1758\log(z)^2-140\log(z)^3-6\log(z)^4\right)}{32},$$

$$\frac{3\left(-17913+23328\,z^{\frac{1}{3}}-6144\,\sqrt{z}+729\,z^{\frac{2}{3}}-5190\ \log(z)-690\log(z)^2-52\log(z)^3-2\log(z)^4\right)}{16}\},$$

$$\{0, 1, \log(z), \frac{\log(z)^2}{2}, \frac{\log(z)^3}{6}, \frac{-17913+23328\,z^{\frac{1}{3}}-6144\,\sqrt{z}+729\,z^{\frac{2}{3}}-5190\ \log(z)-690\log(z)^2-52\log(z)^3}{48},$$

$$\frac{-9\left(-2135+3024\,z^{\frac{1}{3}}-1024\,\sqrt{z}+135\,z^{\frac{2}{3}}-586\ \log(z)-70\log(z)^2-4\log(z)^3\right)}{16},$$

$$\frac{3\left(3\left(-865+1296\,z^{\frac{1}{3}}-512\,\sqrt{z}+81\,z^{\frac{2}{3}}\right)-690\ \log(z)-78\log(z)^2-4\log(z)^3\right)}{8}\},$$

$$\{0, 0, 1, \log(z), \frac{\log(z)^2}{2}, \frac{-865+1296\,z^{\frac{1}{3}}-512\,\sqrt{z}+81\,z^{\frac{2}{3}}-230\ \log(z)-26\log(z)^2}{8},$$

$$\frac{-9\left(-293+504\,z^{\frac{1}{3}}-256\,\sqrt{z}+45\,z^{\frac{2}{3}}-70\ \log(z)-6\log(z)^2\right)}{8}, \frac{9\left(-115+216\,z^{\frac{1}{3}}-128\,\sqrt{z}+27\,z^{\frac{2}{3}}-26\ \log(z)-2\log(z)^2\right)}{4}\},$$

$$\{0, 0, 0, 1, \log(z), \frac{-115+216\,z^{\frac{1}{3}}-128\,\sqrt{z}+27\,z^{\frac{2}{3}}-26\ \log(z)}{4}, \frac{-9\left(-35+84\,z^{\frac{1}{3}}-64\,\sqrt{z}+15\,z^{\frac{2}{3}}-6\ \log(z)\right)}{4},$$

$$\frac{9\left(-13+36\,z^{\frac{1}{3}}-32\,\sqrt{z}+9\,z^{\frac{2}{3}}-2\ \log(z)\right)}{2}\},\ \{0, 0, 0, 0, 1, \frac{-13+36\,z^{\frac{1}{3}}-32\,\sqrt{z}+9\,z^{\frac{2}{3}}}{2}, \frac{-9\left(-3+14\,z^{\frac{1}{3}}-16\,\sqrt{z}+5\,z^{\frac{2}{3}}\right)}{2},$$

$$9\left(-1+z^{\frac{1}{6}}\right)^3\ \left(1+3\,z^{\frac{1}{6}}\right)\},\ \{0, 0, 0, 0, 0, \left(6-8\,z^{\frac{1}{6}}+3\,z^{\frac{1}{3}}\right)\ z^{\frac{1}{3}}, -3\ \left(7-12\,z^{\frac{1}{6}}+5\,z^{\frac{1}{3}}\right)\ z^{\frac{1}{3}},$$

$$18\left(-1+z^{\frac{1}{6}}\right)^2 z^{\frac{1}{3}}\},\ \{0, 0, 0, 0, 0, 2\left(-1+z^{\frac{1}{6}}\right)^2 z^{\frac{1}{3}}, -7\,z^{\frac{1}{3}}+18\,\sqrt{z}-10\,z^{\frac{2}{3}}, 6\ \left(z^{\frac{1}{3}}-3\,\sqrt{z}+2\,z^{\frac{2}{3}}\right)\},$$

$$\{0, 0, 0, 0, 0, \frac{2\left(z^{\frac{1}{3}}-3\,\sqrt{z}+2\,z^{\frac{2}{3}}\right)}{3}, \frac{-\left(\left(7-27\,z^{\frac{1}{6}}+20\,z^{\frac{1}{3}}\right)\,z^{\frac{1}{3}}\right)}{3},\ \left(2-9\,z^{\frac{1}{6}}+8\,z^{\frac{1}{3}}\right)\ z^{\frac{1}{3}}\}\}$$

The matrix q' introduced in (A2) is given by:

$$q' = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & -\left(\frac{391933}{32}\right) & \frac{130077}{32} & -\left(\frac{53739}{16}\right) \\ 0 & 1 & 0 & 0 & 0 & -\left(\frac{5971}{16}\right) & \frac{19215}{16} & -\left(\frac{7785}{8}\right) \\ 0 & 0 & \frac{1}{2} & 0 & 0 & -\left(\frac{865}{16}\right) & \frac{2637}{16} & -\left(\frac{1035}{8}\right) \\ 0 & 0 & 0 & \frac{1}{6} & 0 & -\left(\frac{115}{24}\right) & \frac{105}{8} & -\left(\frac{39}{4}\right) \\ 0 & 0 & 0 & 0 & \frac{1}{24} & -\left(\frac{13}{32}\right) & \frac{9}{16} & -\left(\frac{3}{8}\right) \end{pmatrix} \tag{A7}$$

Further, the matrix q is of the form:

$$q = \begin{pmatrix} q_{00} & q_{01} & q_{02} & q_{03} & q_{04} & q_{05} & q_{06} & q_{07} \\ q_{10} & q_{11} & q_{12} & q_{13} & q_{14} & q_{15} & q_{16} & q_{17} \\ 0 & q_{21} & q_{22} & q_{23} & q_{24} & q_{25} & q_{26} & q_{27} \\ 0 & 0 & 0 & 0 & q_{34} & q_{35} & q_{36} & q_{37} \\ 0 & 0 & 0 & 0 & q_{44} & 0 & 0 & 0 \end{pmatrix} \tag{A8}$$

From the matrix equation $q' = q(M^{-1})^t$, one sees that one has 40 equations in 64 variables, one has the freedom to (judiciously) give arbitrary values to 24 variables, bearing in mind that from the forms of the matrices q' and $q(M^{-1})^t$, the values of $(M^{-1})^t_{4i}$ are fixed. We set: $M_{ij} = \delta_{ij}$ for $0 \leq i \leq 3$ and $0 \leq j \leq 7$.

Hence, one gets:
$q(M^{-1})^t = \{\{q_{00} + (M^{-1})^t_{30}\, q_{03} + (M^{-1})^t_{40}\, q_{04} + (M^{-1})^t_{50}\, q_{05} + (M^{-1})^t_{60}\, q_{06} +$
$(M^{-1})^t_{70}\, q_{07}, q_{01} + (M^{-1})^t_{31}\, q_{03} + (M^{-1})^t_{41}\, q_{04} + (M^{-1})^t_{51}\, q_{05} + (M^{-1})^t_{61}\, q_{06} +$
$(M^{-1})^t_{71}\, q_{07}, q_{02} + (M^{-1})^t_{32}\, q_{03} + (M^{-1})^t_{42}\, q_{04} + (M^{-1})^t_{52}\, q_{05} + (M^{-1})^t_{62}\, q_{06} +$
$(M^{-1})^t_{72}\, q_{07}, (M^{-1})^t_{33}\, q_{03}+(M^{-1})^t_{43}\, q_{04}+(M^{-1})^t_{53}\, q_{05}+(M^{-1})^t_{63}\, q_{06}+(M^{-1})^t_{73}\, q_{07}, (M^{-1})^t_{34}\, q_{03}+$
$(M^{-1})^t_{44}\, q_{04}+(M^{-1})^t_{54}\, q_{05}+(M^{-1})^t_{64}\, q_{06}+(M^{-1})^t_{74}\, q_{07}, (M^{-1})^t_{35}\, q_{03}+(M^{-1})^t_{45}\, q_{04}+$
$(M^{-1})^t_{55}\, q_{05}+(M^{-1})^t_{65}\, q_{06}+(M^{-1})^t_{75}\, q_{07}, (M^{-1})^t_{36}\, q_{03}+(M^{-1})^t_{46}\, q_{04}+(M^{-1})^t_{56}\, q_{05}+$
$(M^{-1})^t_{66}\, q_{06}+(M^{-1})^t_{76}\, q_{07}, (M^{-1})^t_{37}\, q_{03}+(M^{-1})^t_{47}\, q_{04}+(M^{-1})^t_{57}\, q_{05}+(M^{-1})^t_{67}\, q_{06}+$
$(M^{-1})^t_{77}\, q_{07}\}, \{q_{10} + (M^{-1})^t_{30}\, q_{13} + (M^{-1})^t_{40}\, q_{14} + (M^{-1})^t_{50}\, q_{15} + (M^{-1})^t_{60}\, q_{16} +$
$(M^{-1})^t_{70}\, q_{17}, q_{11} + (M^{-1})^t_{31}\, q_{13} + (M^{-1})^t_{41}\, q_{14} + (M^{-1})^t_{51}\, q_{15} + (M^{-1})^t_{61}\, q_{16} +$
$(M^{-1})^t_{71}\, q_{17}, q_{12} + (M^{-1})^t_{32}\, q_{13} + (M^{-1})^t_{42}\, q_{14} + (M^{-1})^t_{52}\, q_{15} + (M^{-1})^t_{62}\, q_{16} +$
$(M^{-1})^t_{72}\, q_{17}, (M^{-1})^t_{33}\, q_{13}+(M^{-1})^t_{43}\, q_{14}+(M^{-1})^t_{53}\, q_{15}+(M^{-1})^t_{63}\, q_{16}+(M^{-1})^t_{73}\, q_{17}, (M^{-1})^t_{34}\, q_{13}+$
$(M^{-1})^t_{44}\, q_{14}+(M^{-1})^t_{54}\, q_{15}+(M^{-1})^t_{64}\, q_{16}+(M^{-1})^t_{74}\, q_{17}, (M^{-1})^t_{35}\, q_{13}+(M^{-1})^t_{45}\, q_{14}+$
$(M^{-1})^t_{55}\, q_{15}+(M^{-1})^t_{65}\, q_{16}+(M^{-1})^t_{75}\, q_{17}, (M^{-1})^t_{36}\, q_{13}+(M^{-1})^t_{46}\, q_{14}+(M^{-1})^t_{56}\, q_{15}+$
$(M^{-1})^t_{66}\, q_{16}+(M^{-1})^t_{76}\, q_{17}, (M^{-1})^t_{37}\, q_{13}+(M^{-1})^t_{47}\, q_{14}+(M^{-1})^t_{57}\, q_{15}+(M^{-1})^t_{67}\, q_{16}+$
$(M^{-1})^t_{77}\, q_{17}\}, \{(M^{-1})^t_{30}\, q_{23}+(M^{-1})^t_{40}\, q_{24}+(M^{-1})^t_{50}\, q_{25}+(M^{-1})^t_{60}\, q_{26}+(M^{-1})^t_{70}\, q_{27}, q_{21}+$
$(M^{-1})^t_{31}\, q_{23} + (M^{-1})^t_{41}\, q_{24} + (M^{-1})^t_{51}\, q_{25} + (M^{-1})^t_{61}\, q_{26} + (M^{-1})^t_{71}\, q_{27}, q_{22} +$
$(M^{-1})^t_{32}\, q_{23}+(M^{-1})^t_{42}\, q_{24}+(M^{-1})^t_{52}\, q_{25}+(M^{-1})^t_{62}\, q_{26}+(M^{-1})^t_{72}\, q_{27}, (M^{-1})^t_{33}\, q_{23}+$
$(M^{-1})^t_{43}\, q_{24}+(M^{-1})^t_{53}\, q_{25}+(M^{-1})^t_{63}\, q_{26}+(M^{-1})^t_{73}\, q_{27}, (M^{-1})^t_{34}\, q_{23}+(M^{-1})^t_{44}\, q_{24}+$
$(M^{-1})^t_{54}\, q_{25}+(M^{-1})^t_{64}\, q_{26}+(M^{-1})^t_{74}\, q_{27}, (M^{-1})^t_{35}\, q_{23}+(M^{-1})^t_{45}\, q_{24}+(M^{-1})^t_{55}\, q_{25}+$
$(M^{-1})^t_{65}\, q_{26}+(M^{-1})^t_{75}\, q_{27}, (M^{-1})^t_{36}\, q_{23}+(M^{-1})^t_{46}\, q_{24}+(M^{-1})^t_{56}\, q_{25}+(M^{-1})^t_{66}\, q_{26}+$
$(M^{-1})^t_{76}\, q_{27}, (M^{-1})^t_{37}\, q_{23}+(M^{-1})^t_{47}\, q_{24}+(M^{-1})^t_{57}\, q_{25}+(M^{-1})^t_{67}\, q_{26}+(M^{-1})^t_{77}\, q_{27}\}, \{(M^{-1})^t_{40}\, q_{34}+$
$(M^{-1})^t_{50}\, q_{35}+(M^{-1})^t_{60}\, q_{36}+(M^{-1})^t_{70}\, q_{37}, (M^{-1})^t_{41}\, q_{34}+(M^{-1})^t_{51}\, q_{35}+(M^{-1})^t_{61}\, q_{36}+$
$(M^{-1})^t_{71}\, q_{37}, (M^{-1})^t_{42}\, q_{34}+(M^{-1})^t_{52}\, q_{35}+(M^{-1})^t_{62}\, q_{36}+(M^{-1})^t_{72}\, q_{37}, (M^{-1})^t_{43}\, q_{34}+$
$(M^{-1})^t_{53}\, q_{35}+(M^{-1})^t_{63}\, q_{36}+(M^{-1})^t_{73}\, q_{37}, (M^{-1})^t_{44}\, q_{34}+(M^{-1})^t_{54}\, q_{35}+(M^{-1})^t_{64}\, q_{36}+$
$(M^{-1})^t_{74}\, q_{37}, (M^{-1})^t_{45}\, q_{34}+(M^{-1})^t_{55}\, q_{35}+(M^{-1})^t_{65}\, q_{36}+(M^{-1})^t_{75}\, q_{37}, (M^{-1})^t_{46}\, q_{34}+$
$(M^{-1})^t_{56}\, q_{35}+(M^{-1})^t_{66}\, q_{36}+(M^{-1})^t_{76}\, q_{37}, (M^{-1})^t_{47}\, q_{34}+(M^{-1})^t_{57}\, q_{35}+(M^{-1})^t_{67}\, q_{36}+$
$(M^{-1})^t_{77}\, q_{37}\}, \{(M^{-1})^t_{40}\, q_{44}, (M^{-1})^t_{41}\, q_{44}, (M^{-1})^t_{42}\, q_{44}, (M^{-1})^t_{43}\, q_{44}, (M^{-1})^t_{44}\, q_{44},$

$(M^{-1})^t_{45}\, q_{44}, (M^{-1})^t_{46}\, q_{44}, (M^{-1})^t_{47}\, q_{44}\}\}$

The above matrix equatin can then be solved for the 64-24=40 entries $(M^{-1})^t_{ij}$ $4 \leq i \leq 7,\ 0 \leq j \leq 7$ to give the following result:

$(M^{-1})^t_{30} = \frac{X_{30}}{Y_{30}}$, where

$X_{30} = q_{06}\, q_{10}\, q_{27}\, q_{35} + q_{16}\, q_{27}\, q_{35} - q_{00}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{10}\, q_{27}\, q_{36} - q_{15}\, q_{27}\, q_{36} +$
$q_{00}\, q_{15}\, q_{27}\, q_{36}(-1 + q_{00})\, q_{17}\, (q_{26}\, q_{35} - q_{25}\, q_{36}) + q_{07}\, q_{10}\, (-(q_{26}\, q_{35}) + q_{25}\, q_{36}) -$
$q_{06}\, q_{10}\, q_{25}\, q_{37} - q_{16}\, q_{25}\, q_{37} + q_{00}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{10}\, q_{26}\, q_{37} + q_{15}\, q_{26}\, q_{37} - q_{00}\, q_{15}\, q_{26}\, q_{37}$

$Y_{30} = -(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} -$
$q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) +$
$q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} +$
$q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}),$

$(M^{-1})^t_{31} = \frac{X_{31}}{Y_{31}}$ where

$X_{31} = q_{01} q_{17} q_{26} q_{35} - q_{01} q_{16} q_{27} q_{35} + q_{05} q_{17} q_{21} q_{36} - q_{01} q_{17} q_{25}\, q_{36} + q_{05} q_{27} q_{36} -$
$q_{05} q_{11} q_{27} q_{36} + q_{01} q_{15} q_{27} q_{36} + q_{07}(q_{16} q_{21} q_{35} + q_{26}(q_{35} - q_{11}\, q_{35}) -$
$(q_{15}\, q_{21} + q_{25} - q_{11}\, q_{25})\, q_{36}) - q_{05}\, q_{16}\, q_{21}\, q_{37} + q_{01}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{26}\, q_{37} + q_{05}\, q_{11}\, q_{26}\, q_{37} -$
$q_{01}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{21}\, q_{35}) + (-1 + q_{11})\, q_{27}\, q_{35} +$
$(q_{15}\, q_{21} + q_{25} - q_{11}\, q_{25})\, q_{37})$

$Y_{31} = -(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} -$
$q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{23}\, q_{37} -$
$q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} +$
$q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}),$

$(M^{-1})^t_{32} = \frac{X_{32}}{Y_{32}}$ where

$X_{32} = 2\, q_{02}\, q_{17}\, q_{26}\, q_{35} - 2\, q_{02}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{36} + 2\, q_{05}\, q_{17}\, q_{22}\, q_{36} - 2\, q_{02}\, q_{17}\, q_{25}\, q_{36} -$
$2\, q_{05}\, q_{12}\, q_{27}\, q_{36} + 2\, q_{02}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, (-1 + 2\, q_{22})\, q_{35} -$
$2\, q_{12}\, q_{26}\, q_{35} + q_{15}\, q_{36} - 2\, q_{15}\, q_{22}\, q_{36} + 2\, q_{12}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{37} - 2\, q_{05}\, q_{16}\, q_{22}\, q_{37} +$
$2\, q_{02}\, q_{16}\, q_{25}\, q_{37} + 2\, q_{05}\, q_{12}\, q_{26}\, q_{37} - 2\, q_{02}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{35} - 2\, q_{17}\, q_{22}\, q_{35} +$
$2\, q_{12}\, q_{27}\, q_{35}$
$-q_{15}\, q_{37} + 2\, q_{15}\, q_{22}\, q_{37} - 2\, q_{12}\, q_{25}\, q_{37})$

$Y_{32} = 2\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36}$
$+q_{05}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} -$
$q_{13}\, q_{25}\, q_{36}) +$
$q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{35} -$
$q_{13}\, q_{27}\, q_{35}$
$-q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37})),$

$(M^{-1})^t_{33} = \frac{X_{33}}{Y_{33}}$ where

$X_{33} = q_{07}\, q_{16}\, q_{25} - q_{06}\, q_{17}\, q_{25} - q_{07}\, q_{15}\, q_{26} + q_{05}\, q_{17}\, q_{26} + q_{06}\, q_{15}\, q_{27} - q_{05}\, q_{16}\, q_{27}$

$Y_{33} = 6\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} +$

$q_{05}\,q_{13}\,q_{27}\,q_{36} - q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(-(q_{16}\,q_{23}\,q_{35}) + q_{13}\,q_{26}\,q_{35}+$
$q_{15}\,q_{23}\,q_{36}-q_{13}\,q_{25}\,q_{36})+q_{05}\,q_{16}\,q_{23}\,q_{37}-q_{03}\,q_{16}\,q_{25}\,q_{37}-q_{05}\,q_{13}\,q_{26}\,q_{37}+q_{03}\,q_{15}\,q_{26}\,q_{37}+$
$q_{06}\,(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37})),$

$(M^{-1})^t_{34} = \frac{X_{34}}{Y_{34}}$ where

$X_{34} = q_{05}\,q_{17}\,q_{26}\,q_{34}-q_{05}\,q_{16}\,q_{27}\,q_{34}-q_{04}\,q_{17}\,q_{26}\,q_{35}+q_{04}\,q_{16}\,q_{27}\,q_{35}-q_{05}\,q_{17}\,q_{24}\,q_{36}+$
$q_{04}\,q_{17}\,q_{25}\,q_{36} + q_{05}\,q_{14}\,q_{27}\,q_{36} - q_{04}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{25}\,q_{34} - q_{15}\,q_{26}\,q_{34}-$
$q_{16}\,q_{24}\,q_{35} + q_{14}\,q_{26}\,q_{35} + q_{15}\,q_{24}\,q_{36} - q_{14}\,q_{25}\,q_{36}) + q_{05}\,q_{16}\,q_{24}\,q_{37} - q_{04}\,q_{16}\,q_{25}\,q_{37} -$
$q_{05}\,q_{14}\,q_{26}\,q_{37}+q_{04}\,q_{15}\,q_{26}\,q_{37}+q_{06}\,(-(q_{17}\,q_{25}\,q_{34})+q_{15}\,q_{27}\,q_{34}+q_{17}\,q_{24}\,q_{35}-q_{14}\,q_{27}\,q_{35}-$
$q_{15}\,q_{24}\,q_{37} + q_{14}\,q_{25}\,q_{37})$

$Y_{34} = 24\,(q_{03}\,q_{17}\,q_{26}\,q_{35}-q_{03}\,q_{16}\,q_{27}\,q_{35}+q_{05}\,q_{17}\,q_{23}\,q_{36}-q_{03}\,q_{17}\,q_{25}\,q_{36}-q_{05}\,q_{13}\,q_{27}\,q_{36}+$
$q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(q_{16}\,q_{23}\,q_{35}-q_{13}\,q_{26}\,q_{35}-q_{15}\,q_{23}\,q_{36}+q_{13}\,q_{25}\,q_{36})-q_{05}\,q_{16}\,q_{23}\,q_{37}+$
$q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37}+$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{35} = \frac{X_{35}}{Y_{35}}$ where

$X_{35} = -39\,q_{05}\,q_{17}\,q_{26}\,q_{34}+39\,q_{05}\,q_{16}\,q_{27}\,q_{34}+39\,q_{04}\,q_{17}\,q_{26}\,q_{35}-39\,q_{04}\,q_{16}\,q_{27}\,q_{35}+$
$39\,q_{05}\,q_{17}\,q_{24}\,q_{36}-39\,q_{04}\,q_{17}\,q_{25}\,q_{36}-39\,q_{05}\,q_{14}\,q_{27}\,q_{36}+39\,q_{04}\,q_{15}\,q_{27}\,q_{36}-39\,q_{05}\,q_{16}\,q_{24}\,q_{37}+$
$39\,q_{04}\,q_{16}\,q_{25}\,q_{37}+39\,q_{05}\,q_{14}\,q_{26}\,q_{37}-39\,q_{04}\,q_{15}\,q_{26}\,q_{37}+460\,q_{05}\,q_{17}\,q_{26}\,q_{44}-460\,q_{05}\,q_{16}\,q_{27}\,q_{44}-$
$1175799\,q_{17}\,q_{26}\,q_{35}\,q_{44}+1175799\,q_{16}\,q_{27}\,q_{35}\,q_{44}-5190\,q_{05}\,q_{17}\,q_{36}\,q_{44}+1175799\,q_{17}\,q_{25}\,q_{36}\,q_{44}+$
$35826\,q_{05}\,q_{27}\,q_{36}\,q_{44}-1175799\,q_{15}\,q_{27}\,q_{36}\,q_{44}+5190\,q_{05}\,q_{16}\,q_{37}\,q_{44}-1175799\,q_{16}\,q_{25}\,q_{37}\,q_{44}-$
$35826\,q_{05}\,q_{26}\,q_{37}\,q_{44} + 1175799\,q_{15}\,q_{26}\,q_{37}\,q_{44} + q_{07}\,(-3\,(q_{26}\,q_{35} - q_{25}\,q_{36})\,(13\,q_{14} -$
$11942\,q_{44})+$
$q_{16}\,(-39\,q_{25}\,q_{34} + 39\,q_{24}\,q_{35} + 460\,q_{25}\,q_{44} - 5190\,q_{35}\,q_{44})+$
$q_{15}\,(39\,q_{26}\,q_{34} - 39\,q_{24}\,q_{36} - 460\,q_{26}\,q_{44} + 5190\,q_{36}\,q_{44}))+$
$q_{06}\,(3\,(q_{27}\,q_{35}-q_{25}\,q_{37})\,(13\,q_{14}-11942\,q_{44})+q_{17}\,(39\,q_{25}\,q_{34}-39\,q_{24}\,q_{35}-460\,q_{25}\,q_{44}+$
$5190\,q_{35}\,q_{44}) + q_{15}\,(-39\,q_{27}\,q_{34} + 39\,q_{24}\,q_{37} + 460\,q_{27}\,q_{44} - 5190\,q_{37}\,q_{44}))$

$Y_{35} = 96\,(q_{03}\,q_{17}\,q_{26}\,q_{35}-q_{03}\,q_{16}\,q_{27}\,q_{35}+q_{05}\,q_{17}\,q_{23}\,q_{36}-q_{03}\,q_{17}\,q_{25}\,q_{36}-q_{05}\,q_{13}\,q_{27}\,q_{36}$
$+q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(q_{16}\,q_{23}\,q_{35}-q_{13}\,q_{26}\,q_{35}-q_{15}\,q_{23}\,q_{36}+q_{13}\,q_{25}\,q_{36})-q_{05}\,q_{16}\,q_{23}\,q_{37}+$
$q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} + q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35}$
$+q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{36} = \frac{X_{36}}{Y_{36}}$ where

$X_{36} = 3\,(6\,q_{05}\,q_{17}\,q_{26}\,q_{34} - 6\,q_{05}\,q_{16}\,q_{27}\,q_{34} - 6\,q_{04}\,q_{17}\,q_{26}\,q_{35} + 6\,q_{04}\,q_{16}\,q_{27}\,q_{35}-$
$6\,q_{05}\,q_{17}\,q_{24}\,q_{36}+6\,q_{04}\,q_{17}\,q_{25}\,q_{36}+6\,q_{05}\,q_{14}\,q_{27}\,q_{36}-6\,q_{04}\,q_{15}\,q_{27}\,q_{36}+6\,q_{05}\,q_{16}\,q_{24}\,q_{37}-$
$6\,q_{04}\,q_{16}\,q_{25}\,q_{37}-6\,q_{05}\,q_{14}\,q_{26}\,q_{37}+6\,q_{04}\,q_{15}\,q_{26}\,q_{37}-140\,q_{05}\,q_{17}\,q_{26}\,q_{44}+140\,q_{05}\,q_{16}\,q_{27}\,q_{44}+$
$43359\,q_{17}\,q_{26}\,q_{35}\,q_{44}-43359\,q_{16}\,q_{27}\,q_{35}\,q_{44}+1758\,q_{05}\,q_{17}\,q_{36}\,q_{44}-43359\,q_{17}\,q_{25}\,q_{36}\,q_{44}-$
$12810\,q_{05}\,q_{27}\,q_{36}\,q_{44}+43359\,q_{15}\,q_{27}\,q_{36}\,q_{44}-1758\,q_{05}\,q_{16}\,q_{37}\,q_{44}+43359\,q_{16}\,q_{25}\,q_{37}\,q_{44}+$
$12810\,q_{05}\,q_{26}\,q_{37}\,q_{44}-43359\,q_{15}\,q_{26}\,q_{37}\,q_{44}+2\,q_{07}\,(3\,(q_{26}\,q_{35}-q_{25}\,q_{36})\,(q_{14}-2135\,q_{44})+$
$q_{16}\,(3\,q_{25}\,q_{34} - 3\,q_{24}\,q_{35} - 70\,q_{25}\,q_{44} + 879\,q_{35}\,q_{44})+$
$q_{15}\,(-3\,q_{26}\,q_{34} + 3\,q_{24}\,q_{36} + 70\,q_{26}\,q_{44} - 879\,q_{36}\,q_{44}))+$
$2\,q_{06}\,(-3\,(q_{27}\,q_{35} - q_{25}\,q_{37})\,(q_{14} - 2135\,q_{44})+$

$q_{17}\,(-3\,q_{25}\,q_{34}+3\,q_{24}\,q_{35}+70\,q_{25}\,q_{44}-879\,q_{35}\,q_{44})+q_{15}\,(3\,q_{27}\,q_{34}-3\,q_{24}\,q_{37}$
$-70\,q_{27}\,q_{44}+879\,q_{37}\,q_{44})))$

$Y_{36}=32\,(q_{03}\,q_{17}\,q_{26}\,q_{35}-q_{03}\,q_{16}\,q_{27}\,q_{35}+q_{05}\,q_{17}\,q_{23}\,q_{36}-q_{03}\,q_{17}\,q_{25}\,q_{36}-q_{05}\,q_{13}\,q_{27}\,q_{36}+$
$q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(q_{16}\,q_{23}\,q_{35}-q_{13}\,q_{26}\,q_{35}-q_{15}\,q_{23}\,q_{36}+q_{13}\,q_{25}\,q_{36})-q_{05}\,q_{16}\,q_{23}\,q_{37}+$
$q_{03}\,q_{16}\,q_{25}\,q_{37}+q_{05}\,q_{13}\,q_{26}\,q_{37}-q_{03}\,q_{15}\,q_{26}\,q_{37}+$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35})+q_{13}\,q_{27}\,q_{35}+q_{15}\,q_{23}\,q_{37}-q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{37}=\frac{X_{37}}{Y_{37}}$ where

$X_{37}=-3\,(2\,q_{05}\,q_{17}\,q_{26}\,q_{34}-2\,q_{05}\,q_{16}\,q_{27}\,q_{34}-2\,q_{04}\,q_{17}\,q_{26}\,q_{35}+2\,q_{04}\,q_{16}\,q_{27}\,q_{35}-$
$2\,q_{05}\,q_{17}\,q_{24}\,q_{36}+2\,q_{04}\,q_{17}\,q_{25}\,q_{36}+2\,q_{05}\,q_{14}\,q_{27}\,q_{36}-2\,q_{04}\,q_{15}\,q_{27}\,q_{36}+2\,q_{05}\,q_{16}\,q_{24}\,q_{37}-$
$2\,q_{04}\,q_{16}\,q_{25}\,q_{37}-2\,q_{05}\,q_{14}\,q_{26}\,q_{37}+2\,q_{04}\,q_{15}\,q_{26}\,q_{37}-52\,q_{05}\,q_{17}\,q_{26}\,q_{44}+52\,q_{05}\,q_{16}\,q_{27}\,q_{44}+$
$17913\,q_{17}\,q_{26}\,q_{35}\,q_{44}-17913\,q_{16}\,q_{27}\,q_{35}\,q_{44}+690\,q_{05}\,q_{17}\,q_{36}\,q_{44}-17913\,q_{17}\,q_{25}\,q_{36}\,q_{44}-$
$5190\,q_{05}\,q_{27}\,q_{36}\,q_{44}+17913\,q_{15}\,q_{27}\,q_{36}\,q_{44}-690\,q_{05}\,q_{16}\,q_{37}\,q_{44}+17913\,q_{16}\,q_{25}\,q_{37}\,q_{44}+$
$5190\,q_{05}\,q_{26}\,q_{37}\,q_{44}-17913\,q_{15}\,q_{26}\,q_{37}\,q_{44}+2\,q_{07}\,((q_{26}\,q_{35}-q_{25}\,q_{36})\,(q_{14}-2595\,q_{44})+$
$q_{16}\,(q_{25}\,q_{34}-q_{24}\,q_{35}-26\,q_{25}\,q_{44}+345\,q_{35}\,q_{44})+$
$q_{15}\,(-(q_{26}\,q_{34})+q_{24}\,q_{36}+26\,q_{26}\,q_{44}-345\,q_{36}\,q_{44}))+2\,q_{06}\,(-((q_{27}\,q_{35}-q_{25}\,q_{37})\,(q_{14}-$
$2595\,q_{44}))$
$+q_{17}\,(-(q_{25}\,q_{34})+q_{24}\,q_{35}+26\,q_{25}\,q_{44}-345\,q_{35}\,q_{44})+$
$q_{15}\,(q_{27}\,q_{34}-q_{24}\,q_{37}-26\,q_{27}\,q_{44}+345\,q_{37}\,q_{44})))$

$Y_{37}=16\,(q_{03}\,q_{17}\,q_{26}\,q_{35}-q_{03}\,q_{16}\,q_{27}\,q_{35}+q_{05}\,q_{17}\,q_{23}\,q_{36}-q_{03}\,q_{17}\,q_{25}\,q_{36}-q_{05}\,q_{13}\,q_{27}\,q_{36}+$
$q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(q_{16}\,q_{23}\,q_{35}-q_{13}\,q_{26}\,q_{35}-q_{15}\,q_{23}\,q_{36}+q_{13}\,q_{25}\,q_{36})-q_{05}\,q_{16}\,q_{23}\,q_{37}+$
$q_{03}\,q_{16}\,q_{25}\,q_{37}+q_{05}\,q_{13}\,q_{26}\,q_{37}-q_{03}\,q_{15}\,q_{26}\,q_{37}+$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35})+q_{13}\,q_{27}\,q_{35}+q_{15}\,q_{23}\,q_{37}-q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{50}=\frac{X_{50}}{Y_{50}}$ where

$X_{50}=-(q_{07}\,q_{10}\,q_{23}\,q_{36})+(-1+q_{00})\,q_{17}\,q_{23}\,q_{36}+q_{03}\,q_{10}\,q_{27}\,q_{36}+q_{13}\,q_{27}\,q_{36}-$
$q_{00}\,q_{13}\,q_{27}\,q_{36}+q_{06}\,q_{10}\,q_{23}\,q_{37}+q_{16}\,q_{23}\,q_{37}-q_{00}\,q_{16}\,q_{23}\,q_{37}-q_{03}\,q_{10}\,q_{26}\,q_{37}-q_{13}\,q_{26}\,q_{37}+$
$q_{00}\,q_{13}\,q_{26}\,q_{37}$

$Y_{50}=-(q_{03}\,q_{17}\,q_{26}\,q_{35})+q_{03}\,q_{16}\,q_{27}\,q_{35}-q_{05}\,q_{17}\,q_{23}\,q_{36}+q_{03}\,q_{17}\,q_{25}\,q_{36}+q_{05}\,q_{13}\,q_{27}\,q_{36}-$
$q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(-(q_{16}\,q_{23}\,q_{35})+q_{13}\,q_{26}\,q_{35}+q_{15}\,q_{23}\,q_{36}-q_{13}\,q_{25}\,q_{36})+q_{05}\,q_{16}\,q_{23}\,q_{37}-$
$q_{03}\,q_{16}\,q_{25}\,q_{37}-q_{05}\,q_{13}\,q_{26}\,q_{37}+q_{03}\,q_{15}\,q_{26}\,q_{37}+q_{06}\,(q_{17}\,q_{23}\,q_{35}-q_{13}\,q_{27}\,q_{35}-$
$q_{15}\,q_{23}\,q_{37}+q_{13}\,q_{25}\,q_{37}),$

$(M^{-1})^t_{51}=\frac{X_{51}}{Y_{51}}$ where

$X_{51}=q_{01}\,q_{17}\,q_{23}\,q_{36}+q_{07}\,(q_{13}\,q_{21}+q_{23}-q_{11}\,q_{23})\,q_{36}-q_{01}\,q_{13}\,q_{27}\,q_{36}-q_{06}\,q_{13}\,q_{21}\,q_{37}-$
$q_{06}\,q_{23}\,q_{37}+q_{06}\,q_{11}\,q_{23}\,q_{37}-q_{01}\,q_{16}\,q_{23}\,q_{37}+q_{01}\,q_{13}\,q_{26}\,q_{37}+q_{03}\,(-(q_{17}\,q_{21}\,q_{36})+$
$(-1+q_{11})\,q_{27}\,q_{36}+(q_{16}\,q_{21}+q_{26}-q_{11}\,q_{26})\,q_{37})$

$Y_{51}=-(q_{03}\,q_{17}\,q_{26}\,q_{35})+q_{03}\,q_{16}\,q_{27}\,q_{35}-q_{05}\,q_{17}\,q_{23}\,q_{36}+q_{03}\,q_{17}\,q_{25}\,q_{36}+q_{05}\,q_{13}\,q_{27}\,q_{36}-$
$q_{03}\,q_{15}\,q_{27}\,q_{36}+q_{07}\,(-(q_{16}\,q_{23}\,q_{35})+q_{13}\,q_{26}\,q_{35}+q_{15}\,q_{23}\,q_{36}-q_{13}\,q_{25}\,q_{36})+q_{05}\,q_{16}\,q_{23}\,q_{37}-$
$q_{03}\,q_{16}\,q_{25}\,q_{37}-q_{05}\,q_{13}\,q_{26}\,q_{37}+q_{03}\,q_{15}\,q_{26}\,q_{37}+q_{06}\,(q_{17}\,q_{23}\,q_{35}-q_{13}\,q_{27}\,q_{35}-$
$q_{15}\,q_{23}\,q_{37}+q_{13}\,q_{25}\,q_{37}),$

$(M^{-1})^t_{52} = \frac{X_{52}}{Y_{52}}$ where

$X_{52} = -2\, q_{02}\, q_{17}\, q_{23}\, q_{36} + q_{07}\, (q_{13} - 2\, q_{13}\, q_{22} + 2\, q_{12}\, q_{23})\, q_{36} + 2\, q_{02}\, q_{13}\, q_{27}\, q_{36} - q_{06}\, q_{13}\, q_{37} + 2\, q_{06}\, q_{13}\, q_{22}\, q_{37} - 2\, q_{06}\, q_{12}\, q_{23}\, q_{37} + 2\, q_{02}\, q_{16}\, q_{23}\, q_{37} - 2\, q_{02}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, (q_{17}\, (-1 + 2\, q_{22})\, q_{36} - 2\, q_{12}\, q_{27}\, q_{36} + q_{16}\, q_{37} - 2\, q_{16}\, q_{22}\, q_{37} + 2\, q_{12}\, q_{26}\, q_{37})$ $Y_{52} = 2\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} + q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} + q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37}))$,

$(M^{-1})^t_{53} = \frac{X_{53}}{Y_{53}}$ where

$X_{53} = -(q_{07}\, q_{16}\, q_{23}) + q_{06}\, q_{17}\, q_{23} + q_{07}\, q_{13}\, q_{26} - q_{03}\, q_{17}\, q_{26} - q_{06}\, q_{13}\, q_{27} + q_{03}\, q_{16}\, q_{27}$

$Y_{53} = 6\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}))$,

$(M^{-1})^t_{54} = \frac{X_{54}}{Y_{54}}$ where

$X_{54} = -(q_{03}\, q_{17}\, q_{26}\, q_{34}) + q_{03}\, q_{16}\, q_{27}\, q_{34} - q_{04}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{24}\, q_{36} + q_{04}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{14}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{34}) + q_{13}\, q_{26}\, q_{34} + q_{14}\, q_{23}\, q_{36} - q_{13}\, q_{24}\, q_{36}) + q_{04}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{24}\, q_{37} - q_{04}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{14}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{34} - q_{13}\, q_{27}\, q_{34} - q_{14}\, q_{23}\, q_{37} + q_{13}\, q_{24}\, q_{37})$

$Y_{54} = 24\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} + q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} + q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37}))\, q_{44}$,

$(M^{-1})^t_{55} = \frac{X_{55}}{Y_{55}}$ where

$X_{55} = 39\, q_{03}\, q_{17}\, q_{26}\, q_{34} - 39\, q_{03}\, q_{16}\, q_{27}\, q_{34} + 39\, q_{04}\, q_{17}\, q_{23}\, q_{36} - 39\, q_{03}\, q_{17}\, q_{24}\, q_{36} - 39\, q_{04}\, q_{13}\, q_{27}\, q_{36} + 39\, q_{03}\, q_{14}\, q_{27}\, q_{36} - 39\, q_{04}\, q_{16}\, q_{23}\, q_{37} + 39\, q_{03}\, q_{16}\, q_{24}\, q_{37} + 39\, q_{04}\, q_{13}\, q_{26}\, q_{37} - 39\, q_{03}\, q_{14}\, q_{26}\, q_{37} - 460\, q_{03}\, q_{17}\, q_{26}\, q_{44} + 460\, q_{03}\, q_{16}\, q_{27}\, q_{44} + 5190\, q_{03}\, q_{17}\, q_{36}\, q_{44} - 1175799\, q_{17}\, q_{23}\, q_{36}\, q_{44} - 35826\, q_{03}\, q_{27}\, q_{36}\, q_{44} + 1175799\, q_{13}\, q_{27}\, q_{36}\, q_{44} - 5190\, q_{03}\, q_{16}\, q_{37}\, q_{44} + 1175799\, q_{16}\, q_{23}\, q_{37}\, q_{44} + 35826\, q_{03}\, q_{26}\, q_{37}\, q_{44} - 1175799\, q_{13}\, q_{26}\, q_{37}\, q_{44} + q_{07}\, (-39\, q_{14}\, q_{23}\, q_{36} + q_{16}\, q_{23}\, (39\, q_{34} - 460\, q_{44}) + 35826\, q_{23}\, q_{36}\, q_{44} + q_{13}\, (-39\, q_{26}\, q_{34} + 39\, q_{24}\, q_{36} + 460\, q_{26}\, q_{44} - 5190\, q_{36}\, q_{44})) + q_{06}\, (3\, q_{23}\, q_{37}\, (13\, q_{14} - 11942\, q_{44}) + q_{17}\, q_{23}\, (-39\, q_{34} + 460\, q_{44}) + q_{13}\, (39\, q_{27}\, q_{34} - 39\, q_{24}\, q_{37} - 460\, q_{27}\, q_{44} + 5190\, q_{37}\, q_{44}))$

$Y_{55} = 96\,(q_{03}\,q_{17}\,q_{26}\,q_{35} - q_{03}\,q_{16}\,q_{27}\,q_{35} + q_{05}\,q_{17}\,q_{23}\,q_{36} - q_{03}\,q_{17}\,q_{25}\,q_{36} - q_{05}\,q_{13}\,q_{27}\,q_{36} +$
$q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}) - q_{05}\,q_{16}\,q_{23}\,q_{37} +$
$q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} +$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{56} = \frac{X_{56}}{Y_{56}}$ where
$X_{56} = 3\,(-6\,q_{03}\,q_{17}\,q_{26}\,q_{34} + 6\,q_{03}\,q_{16}\,q_{27}\,q_{34} - 6\,q_{04}\,q_{17}\,q_{23}\,q_{36} + 6\,q_{03}\,q_{17}\,q_{24}\,q_{36}$
$+6\,q_{04}\,q_{13}\,q_{27}\,q_{36} - 6\,q_{03}\,q_{14}\,q_{27}\,q_{36} + 6\,q_{04}\,q_{16}\,q_{23}\,q_{37} - 6\,q_{03}\,q_{16}\,q_{24}\,q_{37} - 6\,q_{04}\,q_{13}\,q_{26}\,q_{37} +$
$6\,q_{03}\,q_{14}\,q_{26}\,q_{37} + 140\,q_{03}\,q_{17}\,q_{26}\,q_{44} - 140\,q_{03}\,q_{16}\,q_{27}\,q_{44} - 1758\,q_{03}\,q_{17}\,q_{36}\,q_{44} +$
$43359\,q_{17}\,q_{23}\,q_{36}\,q_{44} + 12810\,q_{03}\,q_{27}\,q_{36}\,q_{44} - 43359\,q_{13}\,q_{27}\,q_{36}\,q_{44} + 1758\,q_{03}\,q_{16}\,q_{37}\,q_{44} -$
$43359\,q_{16}\,q_{23}\,q_{37}\,q_{44} - 12810\,q_{03}\,q_{26}\,q_{37}\,q_{44} + 43359\,q_{13}\,q_{26}\,q_{37}\,q_{44} + 2\,q_{07}\,(3\,q_{23}\,q_{36}\,(q_{14} -$
$2135\,q_{44}) + q_{16}\,q_{23}\,(-3\,q_{34} + 70\,q_{44}) +$
$q_{13}\,(3\,q_{26}\,q_{34} - 3\,q_{24}\,q_{36} - 70\,q_{26}\,q_{44} + 879\,q_{36}\,q_{44})) +$
$2\,q_{06}\,(-3\,q_{23}\,q_{37}\,(q_{14} - 2135\,q_{44}) + q_{17}\,q_{23}\,(3\,q_{34} - 70\,q_{44}) +$
$q_{13}\,(-3\,q_{27}\,q_{34} + 3\,q_{24}\,q_{37} + 70\,q_{27}\,q_{44} - 879\,q_{37}\,q_{44})))$
$Y_{56} = 32\,(q_{03}\,q_{17}\,q_{26}\,q_{35} - q_{03}\,q_{16}\,q_{27}\,q_{35} + q_{05}\,q_{17}\,q_{23}\,q_{36} - q_{03}\,q_{17}\,q_{25}\,q_{36} -$
$q_{05}\,q_{13}\,q_{27}\,q_{36} + q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}) -$
$q_{05}\,q_{16}\,q_{23}\,q_{37} + q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} +$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{57} = \frac{X_{57}}{Y_{57}}$ where
$X_{57} = -3\,(-2\,q_{03}\,q_{17}\,q_{26}\,q_{34} + 2\,q_{03}\,q_{16}\,q_{27}\,q_{34} - 2\,q_{04}\,q_{17}\,q_{23}\,q_{36} + 2\,q_{03}\,q_{17}\,q_{24}\,q_{36} +$
$2\,q_{04}\,q_{13}\,q_{27}\,q_{36} - 2\,q_{03}\,q_{14}\,q_{27}\,q_{36} + 2\,q_{04}\,q_{16}\,q_{23}\,q_{37} - 2\,q_{03}\,q_{16}\,q_{24}\,q_{37} - 2\,q_{04}\,q_{13}\,q_{26}\,q_{37} +$
$2\,q_{03}\,q_{14}\,q_{26}\,q_{37} + 52\,q_{03}\,q_{17}\,q_{26}\,q_{44} - 52\,q_{03}\,q_{16}\,q_{27}\,q_{44} - 690\,q_{03}\,q_{17}\,q_{36}\,q_{44} + 17913\,q_{17}\,q_{23}\,q_{36}\,q_{44} +$
$5190\,q_{03}\,q_{27}\,q_{36}\,q_{44} - 17913\,q_{13}\,q_{27}\,q_{36}\,q_{44} + 690\,q_{03}\,q_{16}\,q_{37}\,q_{44} - 17913\,q_{16}\,q_{23}\,q_{37}\,q_{44} -$
$5190\,q_{03}\,q_{26}\,q_{37}\,q_{44} + 17913\,q_{13}\,q_{26}\,q_{37}\,q_{44} + 2\,q_{07}\,(q_{23}\,q_{36}\,(q_{14} - 2595\,q_{44}) - q_{16}\,q_{23}\,(q_{34} -$
$26\,q_{44}) +$
$q_{13}\,(q_{26}\,q_{34} - q_{24}\,q_{36} - 26\,q_{26}\,q_{44} + 345\,q_{36}\,q_{44})) +$
$2\,q_{06}\,(-(q_{23}\,q_{37}\,(q_{14} - 2595\,q_{44})) + q_{17}\,q_{23}\,(q_{34} - 26\,q_{44}) +$
$q_{13}\,(-(q_{27}\,q_{34}) + q_{24}\,q_{37} + 26\,q_{27}\,q_{44} - 345\,q_{37}\,q_{44})))$
$Y_{57} = 16\,(q_{03}\,q_{17}\,q_{26}\,q_{35} - q_{03}\,q_{16}\,q_{27}\,q_{35} + q_{05}\,q_{17}\,q_{23}\,q_{36} - q_{03}\,q_{17}\,q_{25}\,q_{36} - q_{05}\,q_{13}\,q_{27}\,q_{36}$
$+q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}) - q_{05}\,q_{16}\,q_{23}\,q_{37} +$
$q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} +$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{60} = \frac{X_{60}}{Y_{60}}$ where
$X_{60} = -(q_{07}\,q_{10}\,q_{23}\,q_{35}) + (-1 + q_{00})\,q_{17}\,q_{23}\,q_{35} + q_{03}\,q_{10}\,q_{27}\,q_{35} + q_{13}\,q_{27}\,q_{35} -$
$q_{00}\,q_{13}\,q_{27}\,q_{35} + q_{05}\,q_{10}\,q_{23}\,q_{37} + q_{15}\,q_{23}\,q_{37} - q_{00}\,q_{15}\,q_{23}\,q_{37} - q_{03}\,q_{10}\,q_{25}\,q_{37} - q_{13}\,q_{25}\,q_{37} +$
$q_{00}\,q_{13}\,q_{25}\,q_{37}$

$Y_{60} = q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} + q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} + q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37})$,

$(M^{-1})^t_{61} = \frac{X_{61}}{Y_{61}}$ where

$X_{61} = q_{01}\, q_{17}\, q_{23}\, q_{35} + q_{07}\, (q_{13}\, q_{21} + q_{23} - q_{11}\, q_{23})\, q_{35} - q_{01}\, q_{13}\, q_{27}\, q_{35} - q_{05}\, q_{13}\, q_{21}\, q_{37} - q_{05}\, q_{23}\, q_{37} + q_{05}\, q_{11}\, q_{23}\, q_{37} - q_{01}\, q_{15}\, q_{23}\, q_{37} + q_{01}\, q_{13}\, q_{25}\, q_{37} + q_{03}\, (-(q_{17}\, q_{21}\, q_{35}) + (-1 + q_{11})\, q_{27}\, q_{35} + (q_{15}\, q_{21} + q_{25} - q_{11}\, q_{25})\, q_{37})$

$Y_{61} = q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} + q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} + q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37})$,

$(M^{-1})^t_{62} = \frac{X_{62}}{Y_{62}}$ where

$X_{62} = -2\, q_{02}\, q_{17}\, q_{23}\, q_{35} + q_{07}\, (q_{13} - 2\, q_{13}\, q_{22} + 2\, q_{12}\, q_{23})\, q_{35} + 2\, q_{02}\, q_{13}\, q_{27}\, q_{35} - q_{05}\, q_{13}\, q_{37} + 2\, q_{05}\, q_{13}\, q_{22}\, q_{37} - 2\, q_{05}\, q_{12}\, q_{23}\, q_{37} + 2\, q_{02}\, q_{15}\, q_{23}\, q_{37} - 2\, q_{02}\, q_{13}\, q_{25}\, q_{37} + q_{03}\, (q_{17}\, (-1 + 2\, q_{22})\, q_{35} - 2\, q_{12}\, q_{27}\, q_{35} + q_{15}\, q_{37} - 2\, q_{15}\, q_{22}\, q_{37} + 2\, q_{12}\, q_{25}\, q_{37})$

$Y_{62} = 2\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}))$,

$(M^{-1})^t_{63} = \frac{X_{63}}{Y_{63}}$ where

$X_{63} = q_{07}\, q_{15}\, q_{23} - q_{05}\, q_{17}\, q_{23} - q_{07}\, q_{13}\, q_{25} + q_{03}\, q_{17}\, q_{25} + q_{05}\, q_{13}\, q_{27} - q_{03}\, q_{15}\, q_{27}$

$Y_{63} = 6\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}))$,

$(M^{-1})^t_{64} = \frac{X_{64}}{Y_{64}}$ where

$X_{64} = -(q_{03}\, q_{17}\, q_{25}\, q_{34}) + q_{03}\, q_{15}\, q_{27}\, q_{34} - q_{04}\, q_{17}\, q_{23}\, q_{35} + q_{03}\, q_{17}\, q_{24}\, q_{35} + q_{04}\, q_{13}\, q_{27}\, q_{35} - q_{03}\, q_{14}\, q_{27}\, q_{35} + q_{07}\, (-(q_{15}\, q_{23}\, q_{34}) + q_{13}\, q_{25}\, q_{34} + q_{14}\, q_{23}\, q_{35} - q_{13}\, q_{24}\, q_{35}) + q_{04}\, q_{15}\, q_{23}\, q_{37} - q_{03}\, q_{15}\, q_{24}\, q_{37} - q_{04}\, q_{13}\, q_{25}\, q_{37} + q_{03}\, q_{14}\, q_{25}\, q_{37} + q_{05}\, (q_{17}\, q_{23}\, q_{34} - q_{13}\, q_{27}\, q_{34} - q_{14}\, q_{23}\, q_{37} + q_{13}\, q_{24}\, q_{37})$

$Y_{64} = 24\, (-(q_{03}\, q_{17}\, q_{26}\, q_{35}) + q_{03}\, q_{16}\, q_{27}\, q_{35} - q_{05}\, q_{17}\, q_{23}\, q_{36} + q_{03}\, q_{17}\, q_{25}\, q_{36} + q_{05}\, q_{13}\, q_{27}\, q_{36} - q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (-(q_{16}\, q_{23}\, q_{35}) + q_{13}\, q_{26}\, q_{35} + q_{15}\, q_{23}\, q_{36} - q_{13}\, q_{25}\, q_{36}) + q_{05}\, q_{16}\, q_{23}\, q_{37} - q_{03}\, q_{16}\, q_{25}\, q_{37} - q_{05}\, q_{13}\, q_{26}\, q_{37} + q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (q_{17}\, q_{23}\, q_{35} - q_{13}\, q_{27}\, q_{35} - q_{15}\, q_{23}\, q_{37} + q_{13}\, q_{25}\, q_{37}))\, q_{44}$,

$(M^{-1})^t_{65} = \frac{X_{65}}{Y_{65}}$ where

$X_{65} = -39\, q_{03}\, q_{17}\, q_{25}\, q_{34} + 39\, q_{03}\, q_{15}\, q_{27}\, q_{34} - 39\, q_{04}\, q_{17}\, q_{23}\, q_{35} + 39\, q_{03}\, q_{17}\, q_{24}\, q_{35} +$
$39\, q_{04}\, q_{13}\, q_{27}\, q_{35} - 39\, q_{03}\, q_{14}\, q_{27}\, q_{35} + 39\, q_{04}\, q_{15}\, q_{23}\, q_{37} - 39\, q_{03}\, q_{15}\, q_{24}\, q_{37} - 39\, q_{04}\, q_{13}\, q_{25}\, q_{37} +$
$39\, q_{03}\, q_{14}\, q_{25}\, q_{37} + 460\, q_{03}\, q_{17}\, q_{25}\, q_{44} - 460\, q_{03}\, q_{15}\, q_{27}\, q_{44} - 5190\, q_{03}\, q_{17}\, q_{35}\, q_{44} +$
$1175799\, q_{17}\, q_{23}\, q_{35}\, q_{44} + 35826\, q_{03}\, q_{27}\, q_{35}\, q_{44} - 1175799\, q_{13}\, q_{27}\, q_{35}\, q_{44} + 5190\, q_{03}\, q_{15}\, q_{37}\, q_{44} -$
$1175799\, q_{15}\, q_{23}\, q_{37}\, q_{44} - 35826\, q_{03}\, q_{25}\, q_{37}\, q_{44} + 1175799\, q_{13}\, q_{25}\, q_{37}\, q_{44} + q_{07}\, (3\, q_{23}\, q_{35}\, (13\, q_{14} -$
$11942\, q_{44}) +$
$q_{15}\, q_{23}\, (-39\, q_{34} + 460\, q_{44}) + q_{13}\, (39\, q_{25}\, q_{34} - 39\, q_{24}\, q_{35} - 460\, q_{25}\, q_{44} + 5190\, q_{35}\, q_{44})) +$
$q_{05}\, (-39\, q_{14}\, q_{23}\, q_{37} + q_{17}\, q_{23}\, (39\, q_{34} - 460\, q_{44}) + 35826\, q_{23}\, q_{37}\, q_{44} +$
$q_{13}\, (-39\, q_{27}\, q_{34} + 39\, q_{24}\, q_{37} + 460\, q_{27}\, q_{44} - 5190\, q_{37}\, q_{44}))$

$Y_{65} = 96\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} +$
$q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} +$
$q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} +$
$q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37}))\, q_{44},$

$(M^{-1})^t_{66} = \frac{X_{66}}{Y_{66}}$ where

$X_{66} = 3\, (6\, q_{03}\, q_{17}\, q_{25}\, q_{34} - 6\, q_{03}\, q_{15}\, q_{27}\, q_{34} + 6\, q_{04}\, q_{17}\, q_{23}\, q_{35} - 6\, q_{03}\, q_{17}\, q_{24}\, q_{35} -$
$6\, q_{04}\, q_{13}\, q_{27}\, q_{35} + 6\, q_{03}\, q_{14}\, q_{27}\, q_{35} - 6\, q_{04}\, q_{15}\, q_{23}\, q_{37} + 6\, q_{03}\, q_{15}\, q_{24}\, q_{37} + 6\, q_{04}\, q_{13}\, q_{25}\, q_{37} -$
$6\, q_{03}\, q_{14}\, q_{25}\, q_{37} - 140\, q_{03}\, q_{17}\, q_{25}\, q_{44} + 140\, q_{03}\, q_{15}\, q_{27}\, q_{44} + 1758\, q_{03}\, q_{17}\, q_{35}\, q_{44} -$
$43359\, q_{17}\, q_{23}\, q_{35}\, q_{44} - 12810\, q_{03}\, q_{27}\, q_{35}\, q_{44} + 43359\, q_{13}\, q_{27}\, q_{35}\, q_{44} - 1758\, q_{03}\, q_{15}\, q_{37}\, q_{44} +$
$43359\, q_{15}\, q_{23}\, q_{37}\, q_{44} + 12810\, q_{03}\, q_{25}\, q_{37}\, q_{44} - 43359\, q_{13}\, q_{25}\, q_{37}\, q_{44} + 2\, q_{07}\, (-3\, q_{23}\, q_{35}\, (q_{14} -$
$2135\, q_{44}) + q_{15}\, q_{23}\, (3\, q_{34} - 70\, q_{44}) +$
$q_{13}\, (-3\, q_{25}\, q_{34} + 3\, q_{24}\, q_{35} + 70\, q_{25}\, q_{44} - 879\, q_{35}\, q_{44})) + 2\, q_{05}\, (3\, q_{23}\, q_{37}\, (q_{14} - 2135\, q_{44}) +$
$q_{17}\, q_{23}\, (-3\, q_{34} + 70\, q_{44}) + q_{13}\, (3\, q_{27}\, q_{34} - 3\, q_{24}\, q_{37} - 70\, q_{27}\, q_{44} + 879\, q_{37}\, q_{44})))$

$Y_{66} = 32\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36} +$
$q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} +$
$q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} + q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} +$
$q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37}))\, q_{44},$

$(M^{-1})^t_{67} = \frac{X_{67}}{Y_{67}}$ where

$X_{67} = 3\, (-2\, q_{03}\, q_{17}\, q_{25}\, q_{34} + 2\, q_{03}\, q_{15}\, q_{27}\, q_{34} - 2\, q_{04}\, q_{17}\, q_{23}\, q_{35} + 2\, q_{03}\, q_{17}\, q_{24}\, q_{35} +$
$2\, q_{04}\, q_{13}\, q_{27}\, q_{35} - 2\, q_{03}\, q_{14}\, q_{27}\, q_{35} + 2\, q_{04}\, q_{15}\, q_{23}\, q_{37} - 2\, q_{03}\, q_{15}\, q_{24}\, q_{37} - 2\, q_{04}\, q_{13}\, q_{25}\, q_{37} +$
$2\, q_{03}\, q_{14}\, q_{25}\, q_{37} + 52\, q_{03}\, q_{17}\, q_{25}\, q_{44} - 52\, q_{03}\, q_{15}\, q_{27}\, q_{44} - 690\, q_{03}\, q_{17}\, q_{35}\, q_{44} + 17913\, q_{17}\, q_{23}\, q_{35}\, q_{44} +$
$5190\, q_{03}\, q_{27}\, q_{35}\, q_{44} - 17913\, q_{13}\, q_{27}\, q_{35}\, q_{44} + 690\, q_{03}\, q_{15}\, q_{37}\, q_{44} - 17913\, q_{15}\, q_{23}\, q_{37}\, q_{44} -$
$5190\, q_{03}\, q_{25}\, q_{37}\, q_{44} + 17913\, q_{13}\, q_{25}\, q_{37}\, q_{44} + 2\, q_{07}\, (q_{23}\, q_{35}\, (q_{14} - 2595\, q_{44}) - q_{15}\, q_{23}\, (q_{34} -$
$26\, q_{44}) +$
$q_{13}\, (q_{25}\, q_{34} - q_{24}\, q_{35} - 26\, q_{25}\, q_{44} + 345\, q_{35}\, q_{44})) + 2\, q_{05}\, (-(q_{23}\, q_{37}\, (q_{14} - 2595\, q_{44})) +$
$q_{17}\, q_{23}\, (q_{34} - 26\, q_{44}) + q_{13}\, (-(q_{27}\, q_{34}) + q_{24}\, q_{37} + 26\, q_{27}\, q_{44} - 345\, q_{37}\, q_{44})))$

$Y_{67} = 16\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} - q_{05}\, q_{13}\, q_{27}\, q_{36}$
$+ q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) - q_{05}\, q_{16}\, q_{23}\, q_{37} +$
$q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} +$

$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{70} = \frac{X_{70}}{Y_{70}}$ where
$X_{70} = -(q_{23}\,q_{35}\,(-(q_{10}\,(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36}))+$
$(-1 + q_{00})\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}))$
$Y_{70} = (q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36})$
$\times(q_{07}\,q_{23}\,q_{35} - q_{03}\,q_{27}\,q_{35} - q_{05}\,q_{23}\,q_{37} + q_{03}\,q_{25}\,q_{37})-$
$(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36})$
$\times(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37})),$

$(M^{-1})^t_{71} = \frac{X_{71}}{Y_{71}}$ where
$X_{71} = -(-((q_{13}\,q_{21} + q_{23} - q_{11}\,q_{23})\,q_{35}\,(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} +$
$q_{03}\,q_{25}\,q_{36}))$
$+(q_{03}\,q_{21} - q_{01}\,q_{23})\,q_{35}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36})$
$Y_{71} = -((q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36})$
$\times(q_{07}\,q_{23}\,q_{35} - q_{03}\,q_{27}\,q_{35} - q_{05}\,q_{23}\,q_{37} + q_{03}\,q_{25}\,q_{37}))+$
$(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36})$
$\times(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37})),$

$(M^{-1})^t_{72} = \frac{X_{72}}{Y_{72}}$ where
$X_{72} = -(q_{35}\,(-((q_{13}\,(-1{+}2\,q_{22}){-}2\,q_{12}\,q_{23})\,(q_{06}\,q_{23}\,q_{35}{-}q_{03}\,q_{26}\,q_{35}{-}q_{05}\,q_{23}\,q_{36}{+}$
$q_{03}\,q_{25}\,q_{36})) + (q_{03}\,(-1 + 2\,q_{22}) - 2\,q_{02}\,q_{23})\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} +$
$q_{13}\,q_{25}\,q_{36})))$
$Y_{72} = 2\,(-((q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36})\,(q_{07}\,q_{23}\,q_{35} -$
$q_{03}\,q_{27}\,q_{35}-$
$q_{05}\,q_{23}\,q_{37} + q_{03}\,q_{25}\,q_{37})) + (q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36})$
$\times(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37})),$

$(M^{-1})^t_{73} = \frac{X_{73}}{Y_{73}}$ where
$X_{73} = -(q_{06}\,q_{15}\,q_{23}) + q_{05}\,q_{16}\,q_{23} + q_{06}\,q_{13}\,q_{25} - q_{03}\,q_{16}\,q_{25} - q_{05}\,q_{13}\,q_{26} +$
$q_{03}\,q_{15}\,q_{26}$
$Y_{73} = 6\,(-(q_{03}\,q_{17}\,q_{26}\,q_{35}) + q_{03}\,q_{16}\,q_{27}\,q_{35} - q_{05}\,q_{17}\,q_{23}\,q_{36} + q_{03}\,q_{17}\,q_{25}\,q_{36}+$
$q_{05}\,q_{13}\,q_{27}\,q_{36} - q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(-(q_{16}\,q_{23}\,q_{35}) + q_{13}\,q_{26}\,q_{35} + q_{15}\,q_{23}\,q_{36} -$
$q_{13}\,q_{25}\,q_{36}) + q_{05}\,q_{16}\,q_{23}\,q_{37} - q_{03}\,q_{16}\,q_{25}\,q_{37} - q_{05}\,q_{13}\,q_{26}\,q_{37} + q_{03}\,q_{15}\,q_{26}\,q_{37}+$
$q_{06}\,(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37})),$

$(M^{-1})^t_{74} = \frac{X_{74}}{Y_{74}}$ where
$X_{74} = -(-((q_{15}\,q_{23}\,q_{34} - q_{13}\,q_{25}\,q_{34} - q_{14}\,q_{23}\,q_{35} + q_{13}\,q_{24}\,q_{35})$
$\times(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36}))+$
$(q_{05}\,q_{23}\,q_{34} - q_{03}\,q_{25}\,q_{34} - q_{04}\,q_{23}\,q_{35} + q_{03}\,q_{24}\,q_{35})$

$\times(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}))$

$Y_{74} = 24\,(-((q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36})$
$\times(q_{07}\,q_{23}\,q_{35} - q_{03}\,q_{27}\,q_{35} - q_{05}\,q_{23}\,q_{37} + q_{03}\,q_{25}\,q_{37}))+$
$(q_{06}\,q_{23}\,q_{35} - q_{03}\,q_{26}\,q_{35} - q_{05}\,q_{23}\,q_{36} + q_{03}\,q_{25}\,q_{36})$
$\times(q_{17}\,q_{23}\,q_{35} - q_{13}\,q_{27}\,q_{35} - q_{15}\,q_{23}\,q_{37} + q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{75} = \frac{X_{75}}{Y_{75}}$ where

$X_{75} = 39\,q_{03}\,q_{16}\,q_{25}\,q_{34} - 39\,q_{03}\,q_{15}\,q_{26}\,q_{34} + 39\,q_{04}\,q_{16}\,q_{23}\,q_{35} - 39\,q_{03}\,q_{16}\,q_{24}\,q_{35} -$
$39\,q_{04}\,q_{13}\,q_{26}\,q_{35} + 39\,q_{03}\,q_{14}\,q_{26}\,q_{35} - 39\,q_{04}\,q_{15}\,q_{23}\,q_{36} + 39\,q_{03}\,q_{15}\,q_{24}\,q_{36} + 39\,q_{04}\,q_{13}\,q_{25}\,q_{36} -$
$39\,q_{03}\,q_{14}\,q_{25}\,q_{36} - 460\,q_{03}\,q_{16}\,q_{25}\,q_{44} + 460\,q_{03}\,q_{15}\,q_{26}\,q_{44} + 5190\,q_{03}\,q_{16}\,q_{35}\,q_{44} -$
$1175799\,q_{16}\,q_{23}\,q_{35}\,q_{44} - 35826\,q_{03}\,q_{26}\,q_{35}\,q_{44} + 1175799\,q_{13}\,q_{26}\,q_{35}\,q_{44} - 5190\,q_{03}\,q_{15}\,q_{36}\,q_{44} +$
$1175799\,q_{15}\,q_{23}\,q_{36}\,q_{44} + 35826\,q_{03}\,q_{25}\,q_{36}\,q_{44} - 1175799\,q_{13}\,q_{25}\,q_{36}\,q_{44} + q_{06}\,(-39\,q_{14}\,q_{23}\,q_{35} +$
$q_{15}\,q_{23}\,(39\,q_{34} - 460\,q_{44}) + 35826\,q_{23}\,q_{35}\,q_{44} +$
$q_{13}\,(-39\,q_{25}\,q_{34} + 39\,q_{24}\,q_{35} + 460\,q_{25}\,q_{44} - 5190\,q_{35}\,q_{44})) +$
$q_{05}\,(3\,q_{23}\,q_{36}\,(13\,q_{14} - 11942\,q_{44}) + q_{16}\,q_{23}\,(-39\,q_{34} + 460\,q_{44}) +$
$q_{13}\,(39\,q_{26}\,q_{34} - 39\,q_{24}\,q_{36} - 460\,q_{26}\,q_{44} + 5190\,q_{36}\,q_{44}))$

$Y_{75} = 96\,(q_{03}\,q_{17}\,q_{26}\,q_{35} - q_{03}\,q_{16}\,q_{27}\,q_{35} + q_{05}\,q_{17}\,q_{23}\,q_{36} - q_{03}\,q_{17}\,q_{25}\,q_{36} -$
$q_{05}\,q_{13}\,q_{27}\,q_{36} + q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}) -$
$q_{05}\,q_{16}\,q_{23}\,q_{37} + q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} +$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{76} = \frac{X_{76}}{Y_{76}}$ where

$X_{76} = 3\,(-6\,q_{03}\,q_{16}\,q_{25}\,q_{34} + 6\,q_{03}\,q_{15}\,q_{26}\,q_{34} - 6\,q_{04}\,q_{16}\,q_{23}\,q_{35} + 6\,q_{03}\,q_{16}\,q_{24}\,q_{35}$
$+6\,q_{04}\,q_{13}\,q_{26}\,q_{35} - 6\,q_{03}\,q_{14}\,q_{26}\,q_{35} + 6\,q_{04}\,q_{15}\,q_{23}\,q_{36} - 6\,q_{03}\,q_{15}\,q_{24}\,q_{36} - 6\,q_{04}\,q_{13}\,q_{25}\,q_{36} +$
$6\,q_{03}\,q_{14}\,q_{25}\,q_{36} + 140\,q_{03}\,q_{16}\,q_{25}\,q_{44} - 140\,q_{03}\,q_{15}\,q_{26}\,q_{44} - 1758\,q_{03}\,q_{16}\,q_{35}\,q_{44} +$
$43359\,q_{16}\,q_{23}\,q_{35}\,q_{44} + 12810\,q_{03}\,q_{26}\,q_{35}\,q_{44} - 43359\,q_{13}\,q_{26}\,q_{35}\,q_{44} + 1758\,q_{03}\,q_{15}\,q_{36}\,q_{44} -$
$43359\,q_{15}\,q_{23}\,q_{36}\,q_{44} - 12810\,q_{03}\,q_{25}\,q_{36}\,q_{44} + 43359\,q_{13}\,q_{25}\,q_{36}\,q_{44} + 2\,q_{06}\,(3\,q_{23}\,q_{35}\,(q_{14} -$
$2135\,q_{44}) + q_{15}\,q_{23}\,(-3\,q_{34} + 70\,q_{44}) +$
$q_{13}\,(3\,q_{25}\,q_{34} - 3\,q_{24}\,q_{35} - 70\,q_{25}\,q_{44} + 879\,q_{35}\,q_{44})) +$
$2\,q_{05}\,(-3\,q_{23}\,q_{36}\,(q_{14} - 2135\,q_{44}) + q_{16}\,q_{23}\,(3\,q_{34} - 70\,q_{44}) +$
$q_{13}\,(-3\,q_{26}\,q_{34} + 3\,q_{24}\,q_{36} + 70\,q_{26}\,q_{44} - 879\,q_{36}\,q_{44})))$

$Y_{76} = 32\,(q_{03}\,q_{17}\,q_{26}\,q_{35} - q_{03}\,q_{16}\,q_{27}\,q_{35} + q_{05}\,q_{17}\,q_{23}\,q_{36} - q_{03}\,q_{17}\,q_{25}\,q_{36} -$
$q_{05}\,q_{13}\,q_{27}\,q_{36} + q_{03}\,q_{15}\,q_{27}\,q_{36} + q_{07}\,(q_{16}\,q_{23}\,q_{35} - q_{13}\,q_{26}\,q_{35} - q_{15}\,q_{23}\,q_{36} + q_{13}\,q_{25}\,q_{36}) -$
$q_{05}\,q_{16}\,q_{23}\,q_{37} + q_{03}\,q_{16}\,q_{25}\,q_{37} + q_{05}\,q_{13}\,q_{26}\,q_{37} - q_{03}\,q_{15}\,q_{26}\,q_{37} +$
$q_{06}\,(-(q_{17}\,q_{23}\,q_{35}) + q_{13}\,q_{27}\,q_{35} + q_{15}\,q_{23}\,q_{37} - q_{13}\,q_{25}\,q_{37}))\,q_{44},$

$(M^{-1})^t_{77} = \frac{X_{77}}{Y_{77}}$ where

$X_{77} = -3\,(-2\,q_{03}\,q_{16}\,q_{25}\,q_{34} + 2\,q_{03}\,q_{15}\,q_{26}\,q_{34} - 2\,q_{04}\,q_{16}\,q_{23}\,q_{35} + 2\,q_{03}\,q_{16}\,q_{24}\,q_{35} +$
$2\,q_{04}\,q_{13}\,q_{26}\,q_{35} - 2\,q_{03}\,q_{14}\,q_{26}\,q_{35} + 2\,q_{04}\,q_{15}\,q_{23}\,q_{36} - 2\,q_{03}\,q_{15}\,q_{24}\,q_{36} - 2\,q_{04}\,q_{13}\,q_{25}\,q_{36} +$
$2\,q_{03}\,q_{14}\,q_{25}\,q_{36} + 52\,q_{03}\,q_{16}\,q_{25}\,q_{44} - 52\,q_{03}\,q_{15}\,q_{26}\,q_{44} - 690\,q_{03}\,q_{16}\,q_{35}\,q_{44} + 17913\,q_{16}\,q_{23}\,q_{35}\,q_{44} +$

$5190\, q_{03}\, q_{26}\, q_{35}\, q_{44} - 17913\, q_{13}\, q_{26}\, q_{35}\, q_{44} + 690\, q_{03}\, q_{15}\, q_{36}\, q_{44} - 17913\, q_{15}\, q_{23}\, q_{36}\, q_{44} -$
$5190\, q_{03}\, q_{25}\, q_{36}\, q_{44} + 17913\, q_{13}\, q_{25}\, q_{36}\, q_{44} + 2\, q_{06}\, (q_{23}\, q_{35}\, (q_{14} - 2595\, q_{44}) - q_{15}\, q_{23}\, (q_{34} -$
$26\, q_{44}) +$
$q_{13}\, (q_{25}\, q_{34} - q_{24}\, q_{35} - 26\, q_{25}\, q_{44} + 345\, q_{35}\, q_{44})) + 2\, q_{05}\, (-(q_{23}\, q_{36}\, (q_{14} - 2595\, q_{44})) +$
$q_{16}\, q_{23}\, (q_{34} - 26\, q_{44}) + q_{13}\, (-(q_{26}\, q_{34}) + q_{24}\, q_{36} + 26\, q_{26}\, q_{44} - 345\, q_{36}\, q_{44})))$

$Y_{77} = 16\, (q_{03}\, q_{17}\, q_{26}\, q_{35} - q_{03}\, q_{16}\, q_{27}\, q_{35} + q_{05}\, q_{17}\, q_{23}\, q_{36} - q_{03}\, q_{17}\, q_{25}\, q_{36} -$
$q_{05}\, q_{13}\, q_{27}\, q_{36} + q_{03}\, q_{15}\, q_{27}\, q_{36} + q_{07}\, (q_{16}\, q_{23}\, q_{35} - q_{13}\, q_{26}\, q_{35} - q_{15}\, q_{23}\, q_{36} + q_{13}\, q_{25}\, q_{36}) -$
$q_{05}\, q_{16}\, q_{23}\, q_{37} + q_{03}\, q_{16}\, q_{25}\, q_{37} + q_{05}\, q_{13}\, q_{26}\, q_{37} - q_{03}\, q_{15}\, q_{26}\, q_{37} +$
$q_{06}\, (-(q_{17}\, q_{23}\, q_{35}) + q_{13}\, q_{27}\, q_{35} + q_{15}\, q_{23}\, q_{37} - q_{13}\, q_{25}\, q_{37}))\, q_{44},$

$(M^{-1})^t_{40} = (M^{-1})^t_{41} = (M^{-1})^t_{42} = (M^{-1})^t_{43} = 0, (M^{-1})^t_{44} = \frac{1}{24\, q_{44}},$
$(M^{-1})^t_{45} = \frac{-13}{32\, q_{44}}, (M^{-1})^t_{46} = \frac{9}{16\, q_{44}}, (M^{-1})^t_{47} = \frac{-3}{8\, q_{44}}$

In the above expressions for $(M^{-1})^t$, the non-zero $q'_{ij}s,\ 0 \leq i \leq 4,\ 0 \leq j \leq 7$, are given by:

$$
\begin{aligned}
q_{00} &= \frac{8\pi^3}{3}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + i\pi\Big] \\
q_{10} &= \frac{8\pi^3}{3} \\
q_{01} &= 2\pi^2\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + \frac{14}{3}\pi^2\Big] \\
q_{11} &= 4\pi^2 ln\Big(\frac{2^2.3^3}{6^6}\Big) \\
q_{02} &= \frac{4\pi^2}{\sqrt{3}}\Big(\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + \frac{\pi}{\sqrt{3}}\Big]^2 + 5\pi^2\Big) \\
q_{12} &= \frac{8\pi^2}{\sqrt{3}}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + \frac{\pi}{\sqrt{3}}\Big] \\
q_{22} &= \frac{4\pi^2}{\sqrt{3}} \\
q_{03} &= \Big(\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) - \frac{\pi}{\sqrt{3}}\Big]^2 + 5\pi^2\Big) \\
q_{13} &= \frac{8\pi^2}{\sqrt{3}}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) - \frac{\pi}{\sqrt{3}}\Big] \\
q_{23} &= \frac{4\pi 62}{\sqrt{3}} \\
q_{04} &= 4\sqrt{2}\pi\Bigg(\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + i\pi\Big]^4 + \frac{65}{3}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big) + i\pi\Big]^2 + \frac{169\pi^4}{3}
\end{aligned}
$$

$$-1440\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big]+402\zeta(4)\Big]$$

$$q_{14}=4\sqrt{2}\pi\Big[4\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big]^3+\frac{65}{3}-1440\zeta(3)\Big]$$

$$q_{24}=4\sqrt{2}\pi\Big(6\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big]+\frac{65}{3}$$

$$q_{34}=16\sqrt{2}\pi\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big]$$

$$q_{44}=4\sqrt{2}\pi$$

$$q_{05}=2\pi^2\Bigg(\Big[ln\Big(\frac{2^2.33^3}{6^6}\Big)+i\pi\Big)^3+15\pi^2 ln\Big(\frac{2^2.33^3}{6^6}\Big)-356\zeta(3)\Bigg)$$

$$q_{15}=2\pi^2\Bigg(\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big)^2+15\pi^2\Big]\Bigg)$$

$$q_{25}=6\pi^2\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big]\Bigg)$$

$$q_{35}=2\pi^2$$

$$q_{06}=\frac{4\pi^2}{\sqrt{3}}\Bigg(\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi+\frac{\pi}{\sqrt{3}}\Big]^3+16\pi^2\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi+\frac{\pi}{\sqrt{3}}\Big]$$

$$-360\zeta(3)-\frac{8\pi^3}{3\sqrt{3}}\Bigg)$$

$$q_{16}=\frac{4\pi^2}{\sqrt{3}}\Bigg(3\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi+\frac{\pi}{\sqrt{3}}\Big]^2+16\pi^2\Bigg)$$

$$q_{26}=\frac{4\pi^2}{\sqrt{3}}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)+i\pi\Big)$$

$$q_{36}=\frac{4\pi^2}{\sqrt{3}}$$

$$q_{07}=\frac{4\pi^2}{\sqrt{3}}\Bigg(\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi-\frac{\pi}{\sqrt{3}}\Big]^3+16\pi^2\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi-\frac{\pi}{\sqrt{3}}\Big]$$

$$-360\zeta(3)+\frac{8\pi^3}{3\sqrt{3}}\Bigg)$$

$$q_{17}=\frac{4\pi^2}{\sqrt{3}}\Bigg(3\Big[ln\Big(\frac{2^2.3^3}{6^6}+i\pi-\frac{\pi}{\sqrt{3}}\Big]^2+16\pi^2\Bigg)$$

$$q_{27}=\frac{4\pi^2}{\sqrt{3}}\Big[ln\Big(\frac{2^2.3^3}{6^6}\Big)-i\pi\Big)$$

$$q_{36} = \frac{4\pi^2}{\sqrt{3}} \tag{A9}$$

The matrix $(M^{-1})^t$ is non-singular as the determinant is non-zero. [11]. From the above the matrix M is found to be as given in **2**.

B Monodromy around $z = \infty$

In this appendix we give the details on the evaluation of the monodromy matrix around $z \to \infty$. The non-zero entries of the 8×5 matrix $A_{ai}(\infty)$ with $a = 0, ..., 7$ $i = 1, ..., 5$ for $z \to \infty$, are given below:

$$A_{01}(\infty) = -\frac{36\prod_{j=1}^{5}\Gamma(\frac{j}{6})\Gamma(\frac{2}{3})\Gamma(\frac{1}{2})}{\Gamma(\frac{7}{6})(\Gamma(\frac{5}{6}))^2} = -(2\pi)^{\frac{5}{2}}6^{\frac{3}{2}}\frac{\sqrt{\pi}\Gamma(\frac{2}{3})}{\Gamma(\frac{7}{6})(\Gamma(\frac{5}{6})^2}$$

$$A_{02}(\infty) = 54\sqrt{3}\pi^{\frac{3}{2}}\frac{\Gamma(\frac{5}{6})}{\Gamma(\frac{4}{3})\Gamma(\frac{2}{3})}$$

$$A_{03}(\infty) = -\frac{4\Gamma(\frac{5}{6}\Gamma(\frac{7}{6}\Gamma(-\frac{1}{3})\Gamma(-\frac{1}{6})\Gamma(\frac{1}{6})\Gamma(\frac{1}{3})}{\Gamma(\frac{3}{2})(\Gamma(\frac{1}{2})^2} = -288\sqrt{\frac{\pi}{3}}$$

$$A_{04}(\infty) = -\frac{9}{4}\frac{\Gamma(\frac{7}{6})\Gamma(\frac{3}{2})\Gamma(\frac{4}{3})\Gamma(-\frac{1}{2})\Gamma(-\frac{1}{3})\Gamma(-\frac{1}{6})\Gamma(\frac{1}{6})}{\Gamma(\frac{5}{3})(\Gamma(\frac{1}{3}))^2} = -3\pi^2\frac{\Gamma(\frac{7}{6})\Gamma(-\frac{1}{3})}{\Gamma(\frac{5}{3})\Gamma(\frac{1}{3})}$$

$$A_{05}(\infty) = -\frac{36}{25}\frac{\Gamma(\frac{4}{3})\Gamma(\frac{7}{6})\Gamma(\frac{3}{2})\Gamma(-\frac{2}{3})\Gamma(-\frac{1}{2})\Gamma(-\frac{1}{6})\Gamma(-\frac{1}{3})}{\Gamma(\frac{11}{6})(\Gamma(\frac{1}{6})^2} = \frac{12\pi}{25\sqrt{3}}\frac{\Gamma(-\frac{2}{3})\Gamma(-\frac{1}{6})}{\Gamma(\frac{11}{6})\Gamma(\frac{1}{6})};$$

$$A_{11}(\infty) = \frac{216(\Gamma(\frac{2}{3})^2\Gamma(\frac{1}{3})}{(\Gamma(\frac{5}{6}))^2}$$

$$A_{13}(\infty) = \frac{8\Gamma(-\frac{1}{3}\Gamma(\frac{1}{3})}{(\Gamma(\frac{1}{2})^2} = -\frac{16}{\sqrt{3}\pi}$$

$$A_{15}(\infty) = \frac{216}{125}\frac{\Gamma(\frac{4}{3}\Gamma(-\frac{2}{3})\Gamma(-\frac{1}{3}}{(\Gamma(\frac{1}{6})^2} = -\frac{144\sqrt{3}\pi}{125}\frac{\Gamma(-\frac{2}{3})}{(\Gamma(\frac{1}{6}))^2};$$

---

[11] The determinant is given by: $\frac{P}{Q}$ where $P = \frac{10613001429\,i}{131072}$ and $Q = \sqrt{2}\,\pi^7\,(8\,(-7\,i + 325\,\sqrt{3})\,\pi^7 + 104\,i\,(87\,i + 35\,\sqrt{3})\,\pi^6\,\log(432) - 2\,\pi^5\,(3423 - 889\,i\,\sqrt{3} + 62208\,(-i + \sqrt{3})\,\log\,\log(432) - 6\,(-742\,i + 393\,\sqrt{3})\,\log(432)^2) - 108\,i\,\log(432)^2\,\zeta(3) + 6\,\pi^4\,((-1351\,i + 53\,\sqrt{3})\,\log(432) + 4478976\,i\,(3\,i + \sqrt{3})\,\log^2\,\log(432) - 20736\,i\,\sqrt{3}\,\log\,\log(432)^2 + 24\,i\,(9\,i + 28\,\sqrt{3})\,\log(432)^3 + 16\,(-3 + 7\,i\,\sqrt{3})\,\zeta(3)) + 9\,i\,\pi\,\log(432)\,(3\,\sqrt{3}\log(432)^3 - 8\,(-3\,i + \sqrt{3})\,\zeta(3)) + 3\,\pi^3\,\log(432)\,((-495 + 2\,i\,\sqrt{3})\,\log(432) - 26873856\,i\,\log^2\,\log(432) - 180\,i\,\log(432)^3 + 96\,(-3\,i + \sqrt{3})\,\zeta(3)) + 36\,\pi^2\,(3\,(-5\,i + \sqrt{3})\,\log(432)^3 - 2\,(23\,i + \sqrt{3})\,\zeta(3) + 20\,i\,\sqrt{3}\log(432)^2\,\zeta(3)))$

$$
\begin{aligned}
A_{21}(\infty) &= \frac{216(\Gamma(\tfrac{1}{2}))^2\Gamma(\tfrac{2}{3})}{(\Gamma(\tfrac{5}{6}))^2} = \frac{216\pi\Gamma(\tfrac{2}{3})}{(\Gamma(\tfrac{5}{6}))^2} \\
A_{24}(\infty) &= \frac{27\Gamma(-\tfrac{1}{2}\Gamma(\tfrac{1}{6})}{8(\Gamma(\tfrac{1}{3})^2} \\
A_{25}(\infty) &= \frac{216}{125}\frac{\Gamma(\tfrac{7}{6})\Gamma(-\tfrac{2}{3})\Gamma(-\tfrac{1}{6})}{\Gamma(\tfrac{1}{6})} = \frac{36}{125}\Gamma(-\frac{2}{3})\Gamma(-\frac{1}{6}) \\
A_{31}(\infty) &= \frac{216\Gamma(\tfrac{1}{6})\Gamma(\tfrac{2}{3})}{\Gamma(\tfrac{5}{6})} \\
A_{32}(\infty) &= \frac{27\Gamma(-\tfrac{1}{6})\Gamma(-\tfrac{1}{2})}{(\Gamma(\tfrac{2}{3})^2} = \frac{27\sqrt{\pi}\Gamma(-\tfrac{1}{6})}{(\Gamma(\tfrac{2}{3}))^2} \\
A_{32}(\infty) &= \frac{27}{8}\frac{\Gamma(\tfrac{3}{2})\Gamma(-\tfrac{2}{3})\Gamma(-\tfrac{1}{2})}{(\Gamma(\tfrac{1}{6}))^2} = -\frac{27\pi}{8}\frac{\Gamma(-\tfrac{2}{3})}{(\Gamma(\tfrac{1}{6}))^2}; \\
A_{41}(\infty) &= -216(\Gamma(\frac{1}{6}))^2\Gamma(\frac{2}{3}) \\
A_{45}(\infty) &= -\frac{216}{125}(\Gamma(\frac{5}{6}))^2\Gamma(-\frac{2}{3}); \\
A_{51}(\infty) &= -216\frac{\Gamma(\tfrac{1}{6})\Gamma(\tfrac{1}{3})\Gamma(\tfrac{2}{3})}{\Gamma(\tfrac{5}{6})} = -\frac{432\pi\Gamma(\tfrac{1}{6})}{\sqrt{3}\Gamma(\tfrac{5}{6})} \\
A_{53}(\infty) &= -8\Gamma(-\frac{1}{3})\Gamma(\frac{1}{3}) = 2\sqrt{3}\pi \\
A_{55}(\infty) &= -\frac{216}{125}\frac{\Gamma(\tfrac{5}{6})\Gamma(-\tfrac{2}{3})\Gamma(-\tfrac{1}{3})}{\Gamma(\tfrac{1}{6})}; \\
A_{61}(\infty) &= \frac{216\Gamma(\tfrac{1}{6})(\Gamma(\tfrac{1}{2}))^2\Gamma(\tfrac{2}{3})}{\Gamma(\tfrac{5}{6})} = \frac{216\pi\Gamma(\tfrac{1}{6})\Gamma(\tfrac{2}{3})}{\Gamma(\tfrac{5}{6})} \\
A_{64}(\infty) &= \frac{27}{8}\frac{\Gamma(\tfrac{2}{3}\Gamma(\tfrac{1}{6})\Gamma(-\tfrac{1}{2})}{\Gamma(\tfrac{1}{3})} = -\frac{27\sqrt{\pi}}{4}\frac{\Gamma(\tfrac{2}{3})\Gamma(\tfrac{1}{6})}{\Gamma(\tfrac{1}{3})} \\
A_{65}(\infty) &= \frac{216}{125}\frac{\Gamma(\tfrac{5}{6})\Gamma(\tfrac{7}{6}\Gamma(-\tfrac{2}{3})\Gamma(-\tfrac{1}{6})}{\Gamma(\tfrac{1}{6})}; \\
A_{71}(\infty) &= 216(\Gamma(\frac{1}{6}))^2\Gamma(\frac{2}{3}) \\
A_{72}(\infty) &= 27\Gamma(\frac{1}{3})\Gamma(-\frac{1}{6})\Gamma(\frac{1}{2}) = 27\sqrt{\pi}\Gamma(\frac{1}{3})\Gamma(-\frac{1}{6}) \\
A_{75}(\infty) &= \frac{216}{125}\frac{\Gamma(\tfrac{5}{6}\Gamma(\tfrac{3}{2})\Gamma(-\tfrac{2}{3})\Gamma(-\tfrac{1}{2})}{\Gamma(\tfrac{1}{6})} = -\frac{216}{125}\frac{\Gamma(\tfrac{5}{6})\Gamma(-\tfrac{2}{3})}{\Gamma(\tfrac{1}{6})}.
\end{aligned}
\tag{B1}
$$

Given that $(T_u)_{ij} = e^{-\frac{\sqrt{-1}\pi i}{3}}\delta_{ij}$, no sum over i, one sees that the equation (41) becomes:

$$e^{-\frac{\sqrt{-1}\pi j}{3}}A_{aj}(\infty) = T_{ab}(\infty)A_{bj}(\infty), \tag{B2}$$

(no sum over j) which needs to be solved for $T_{ab}(\infty)$. MATHEMATICA is unable to perform the required computation - however, it is in principle, doable.

References

[1] K. Hori and C. Vafa, *Mirror symmetry*, arXiv:hep-th/0002222.

[2] A. Misra, *On (orientifold of) type IIA on a compact Calabi-Yau*, Fortsch. der Physik, **52** (2004), arXiv:hep-th/0304209.

[3] A. Misra, *An N = 1 triality by spectrum matching*, to appear in Int.Jour. Mod. Phys. A, arXiv:hep-th/0212054.

[4] M. Bodner, A. C. Cadavid and S. Ferrara, *(2,2) Vacuum Configurations For Type Iia Superstrings: N=2 Supergravity Lagrangians And Algebraic Geometry*, Class. Quant. Grav. **8** (1991) 789.

[5] C. Vafa and E. Witten, *Dual string pairs with N = 1 and N = 2 supersymmetry in four dimensions,* Nucl. Phys. Proc. Suppl. **46**, 225 (1996) [arXiv:hep-th/9507050].

[6] S. Kachru and C. Vafa, *Exact results for N=2 compactifications of heterotic strings*, Nucl. Phys. B **450**, 69 (1995) [arXiv:hep-th/9505105].

[7] E. Witten, *Phases of N = 2 theories in two dimensions*, Nucl. Phys. B **403**, 159 (1993) [arXiv:hep-th/9301042].

[8] B. R. Greene and C. I. Lazaroiu, *Collapsing D-branes in Calabi-Yau moduli space. I,* Nucl. Phys. B **604**, 181 (2001) [arXiv:hep-th/0001025]

[9] S. Hosono, A. Klemm, S. Theisen and S. T. Yau, *Mirror symmetry, mirror map and applications to Calabi-Yau hypersurfaces*, Commun. Math. Phys. **167** (1995) 301 [arXiv:hep-th/9308122].

[10] D. R. Morrison, *Picard-Fuchs equations and mirror maps for hypersurfaces,* arXiv:hep-th/9111025.

[11] E. Witten, *Non-Perturbative Superpotentials In String Theory*, Nucl. Phys. B **474**, 343 (1996) [arXiv:hep-th/9604030].

[12] D. E. Diaconescu, B. Florea and A. Misra, *Orientifolds, unoriented instantons and localization*, JHEP **0307**, 041 (2003) [arXiv:hep-th/0305021].

[13] C. Vafa, *Topological Landau-Ginzburg Models*, Mod. Phys. Lett. A **6**, 337 (1991).

[14] S. Kachru, M. B. Schulz and S. Trivedi, *Moduli stabilization from fluxes in a simple IIB orientifold*, [arXiv:hep-th/0201028].

[15] J. Louis, J. Sonnenschein, S. Theisen and S. Yankielowicz, *Non-perturbative properties of heterotic string vacua compactified on* $K3\times T^2$, Nucl. Phys. B **480**, 185 (1996) [arXiv:hep-th/9606049].

[16] G. L. Cardoso, G. Curio, D. Lust and T. Mohaupt, *Instanton Numbers and Exchange Symmetries in* $N=2$ *Dual String Pairs*, Phys. Lett. B **382**, 241 (1996) [arXiv:hep-th/9603108].].

[17] G. Aldazabal, A. Font, L. E. Ibanez and F. Quevedo, *'Chains of N=2, D=4 heterotic/type II duals*, Nucl. Phys. B **461**, 85 (1996) [arXiv:hep-th/9510093].

[18] G.L.Cardoso, G.Curio, G.Dall'Agata, D.Lüst, P.Manousselis and G.Zoupanos, *Non-Kähler Backgrounds and Their Five Torsion Classes*, Nucl. Phys. **B652** (2003) 5 [hep-th/0211118]; S.Gueieri, J.Louis, Micu and D.Waldram, *Mirror Symmetry in Generalized Calabi-Yau Compactifications*, Nucl. Phys. **B654** (203), 61 [hep-th/0211102]; K.Becker, M.becker, K.Dasgupta and P.S.Green. *Compactifications of Heterotic Theory on Non-Kähler Complex Manifolds: I*, JHEP 0304(2003) 007 [hep-th/0301161]; K.becker and K.Dasgupta, *Heterotic Strings with Torsion*, JHEP 0211 (2002) 006[hep-th/0209077].